

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 April 2002 (11.04.2002)

PCT

(10) International Publication Number
WO 02/28829 A2

- (51) International Patent Classification⁷: **C07D 207/00**
- (21) International Application Number: **PCT/US01/29926**
- (22) International Filing Date:
24 September 2001 (24.09.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/234,967 25 September 2000 (25.09.2000) US
09/761,850 18 January 2001 (18.01.2001) US
- (71) Applicant: **QUESTCOR PHARMACEUTICALS, INC.**
[US/US]; 3260 Whipple Road, Union City, CA 94587-1217 (US).
- (72) Inventors: **CHONG, Lee**; 37469 Marsten Drive, Nemark, CA 94560 (US). **FRECHETTE, Roger**; 40 Estate Lane, Reading, MA 01867 (US). **SCOTT, Carole**; 6593 Flanders Drive, Newark, CA 94560 (US). **TESTER, Richard**; 877 Heatherstone Way, Mountain View, CA 94040 (US). **SMITH, Whitney**; 1122 Richmond Street, El Cerrito, CA 94530 (US). **CHIBA, Katsumi**; Enoki 33-94 Suita, Osaka 564-0053 (JP). **SAKAMOTO, Masatoshi**; 2-11-6-706 Miyahara Yodogawa, Osaka 532-0003 (JP). **GLUCHOWSKI, Charles**; 154 Coolspring Court, Danville, CA 94506 (US).
- (74) Agents: **POISSANT, Brian, M.** et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/28829 A2

(54) Title: **PEPTIDE DEFORMYLASE INHIBITORS**

(57) Abstract: The invention is directed to a novel class of compounds, which inhibits peptide deformylase, pharmaceutical compositions containing compounds that inhibit peptide deformylase, and method of treating various infections.

PEPTIDE DEFORMYLASE
INHIBITORS

1. FIELD OF THE INVENTION

5 The present invention relates to a novel class of compounds, which inhibit peptide deformylase, pharmaceutical compositions containing compounds that inhibit peptide deformylase, and methods of treating various infections.

2. BACKGROUND OF THE INVENTION

10 One of the most grave problems facing mankind is the continual battle against infectious diseases. Infectious diseases are the leading cause of death worldwide and the third leading cause of death in the United States. A particularly troublesome trend that has emerged in recent years is that infectious diseases that were once susceptible to treatment with antibiotics are now becoming resistant to those antibiotics. Consider, for example, the
15 bacterium *Staphylococcus aureus*, the most common cause of hospital-acquired infection. The discovery that penicillin killed bacteria led to the common use of penicillin during World War II in army hospitals. Soon after the debut of penicillin, strains of *Staphylococcus aureus* were isolated that were resistant to penicillin. In fact, these resistant strains were commonly found in hospitals where penicillin was heavily used. The
20 chemically modified penicillin derivative methicillin was developed and found to be effective against penicillin resistant strains, but incredibly, soon after the introduction of methicillin, new methicillin resistant strains were isolated. Currently, many methicillin resistant *Staphylococcus aureus* strains have been characterized and are causing significant issues both within and outside the hospital setting. It appears that the capability of bacteria
25 to become resistant to drugs is weakening our ability to fight bacterial infections.

 New antibiotics are desperately needed as clinically significant bacterial pathogens have acquired resistance to nearly all existing antibiotics (Chopra et al., 1997, *Antimicrob. Agents Chemother.*, 37:1563-1571; Cohen, 1992, *Science*, 257:1050-1055; Kunin, 1993, *Ann. Intern. Med.*, 118:557-561; Neu, 1992, *Science*, 257:1064-1073; Tenover and Hughes,
30 1996, *JAMA*, 275:300-304).

 Bacteria acquire resistance to antimicrobial drugs through a remarkable variety of mechanisms (Russell and Chopra, 1996, *Understanding antibacterial action and resistance*, 2nd ed. Ellis Horwood, New York, NY; Jacobs, 1994, *Clin. Infect. Dis.*, 19:1-10; Hooper and Wolfson, 1993, *Am. Soc. Microbiol.*, 1993:97-118). The robust fitness of
35 microorganisms is a manifestation of their short replication times and their ability to evolve

in the face of the selective pressure exerted by antibiotics. In order to survive, microorganisms have developed the ability to adapt quickly and effectively to changes in the environment such as changes in the light intensity, oxygen levels, acidity and exposure to antibiotics. The emergence and spread of resistant bacteria is primarily caused by acquisition of drug resistance genes resulting in a broad spectrum of antibiotic resistance (e.g., extended-spectrum cephalosporin-resistant mutant beta-lactamases found in several bacterial organisms). Genetic exchange of multiple-resistance genes, by transformation, transduction and conjugation, combined with selective pressures in settings such as hospitals where there is heavy use of antibiotic therapies, has enhanced the survival and proliferation of antimicrobial agent-resistant bacterial strains occurring by, e.g., spontaneous mutations. *Id.* Resistance has inevitably developed to all antimicrobial agents, although the extent to which bacteria develop resistance to antimicrobial drugs and the speed with which they do so varies with type. (Gold and Moellering, Jr., 1996, *New Eng. J. Med.*, 335(19):1445-1453). Prevention of life threatening microbial infections coupled with medical practice aimed at minimizing the development of drug resistance are certainly important (Moellering, 1990, *Scand. J. Infect. Dis., Suppl.* 70:18-24; McCaig and Hughes, 1995, *JAMA* 273:214-219; Guillemot, 1998, *JAMA*, 279:365-370), but to effectively battle infectious diseases, it is necessary to develop new antimicrobial drugs.

The biochemical apparatus for the synthesis of polypeptides in bacteria and eukaryotes is strikingly different. It is preferable to have antimicrobial drugs that disrupt bacterial metabolism but that do not interfere with the host's metabolism. Antimicrobial drugs that interfere with the biochemical machinery of the host may cause the deleterious side effects associated with certain drugs. Thus, there is a need for antimicrobial drugs that specifically inhibit bacterial metabolism while not effecting eukaryotic biochemistry.

A significant difference between bacterial and eukaryotic protein synthesis is the use of formylated methionine as the first amino acid used during ribosome based synthesis of polypeptides. In most cases eukaryotic protein biosynthesis is initiated with methionine whereas bacterial biosynthesis commences with N-formyl methionine. Proper protein folding and function in bacteria usually occurs only after the N-formyl group is removed from the nascent polypeptide chain. This is accomplished by an enzyme called peptide deformylase (PDF).

A variety of hydroxamic acid derivatives have been previously disclosed. One of the first examples discovered was actinonin. After the discovery of actinonin, and the deconstruction that these types of compounds inhibit bacterial growth it was found that they inhibit matrix metalloproteases. For relevant literature see e.g., *J. Chem Soc. Perk. Tran.*


1(9) 819-860, WO 98/18754, WO 99/57097, US Patent Nos. 5,712,300, 4,325,964, 4,225,617.

WO 99/59568 describes methods of using certain hydroxamic acid derivatives for antibacterial compositions. Large genera of compounds are disclosed yet only a few
5 examples are given and the biological activities are not reported. It is not even known if the compounds disclosed therein are selective inhibitors of peptide deformylase.


There is a need to develop new antimicrobial drugs in view of the capability of microbes to develop resistance to existing drugs; in particular there is a need for drugs that specifically affect the micro-organisms and not the host. The present invention is meant to
10 address these and other needs.

3. SUMMARY OF THE INVENTION


The present invention relates to compounds represented by formulas I-IV:




Formula I



Formula II



Formula III



Formula IV

or a pharmaceutically acceptable salt of such compounds, wherein:

Z is NHOH or OR_a wherein R_a is H, alkyl or a biocleavable moiety;

X is C=O or O=S=O;

Y is heteroalkyl, heterocyclic, or substituted derivatives thereof;

R₁ is C₁-C₁₀ alkyl, aryl, heterocyclic, heteroalkyl or a substituted derivative thereof;

R₂ and R₃ together with nitrogen represent a 4, 5, 6, or 7 membered heterocyclic ring optionally substituted with one of the following substituents -OH, -CH₂OH,

-O-C(=O)-heterocyclic, -O-C(=O)-NH-aryl, -NH-C(=O)-aryl,

-C(=O)-N(C₂-C₆ alkyl)₂, -NH-(C₁-C₆ alkyl)-heterocyclic,

-NH-C(=O)-NH-alkyl, -NH-C(=O)-NH-aryl, -NH-C(=O)-NH-heterocyclic;

or

R₂ and R₄ together with nitrogen form a ring through a -CH₂-CH₂- linkage wherein

R₃ is H, C₁-C₁₀ alkyl, aryl, heterocyclic, heteroalkyl or a substituted

derivative thereof; or R₂ is methyl;

R₃, as defined above forms a ring with R₂, or represents H, or is alkyl, heteroalkyl, aryl, heterocyclic, and substituted derivatives thereof;

R₃ when not in a ring with R₂ is alkyl, heteroalkyl, aryl, heterocyclic, and substituted derivatives thereof;

R₄ when not in a ring with R₂ is H, C₁-C₁₀ alkyl, aryl, heterocyclic, heteroalkyl or a substituted derivative thereof; or is together with R₂ and nitrogen as defined above to form a ring through a -CH₂-CH₂- linkage;

R₅ is -H, -NO₂, -NH₂, -NH-C(=O)-H, -NH-C(=O)-CH₃, -NH-S(O₂)-CH₃,

-CH₂-NH-alkyl, -CH₂-NH-heteroalkyl, -CH₂-NH-heterocycl, or substituted derivatives thereof;

R₆ is -H, -NO₂, -NH₂, -NH-C(=O)-H, -NH-C(=O)-CH₃, -NH-S(O₂)-CH₃,

-CH₂-NH-alkyl, -CH₂-NH-heteroalkyl, -CH₂-NH-heterocycl, or substituted derivatives thereof;

R_7 or R_8 is $-\text{CHR}_{10}-\text{C}(=\text{O})-\text{NH}-\text{OH}$; R_7 or R_8 when not $-\text{CHR}-\text{C}(=\text{O})-\text{NH}-\text{OH}$ is alkyl, heteroalkyl, heterocycyl, alkylaryl, alkylheterocyclic, or substituted derivatives thereof;

5 R_9 is H, alkyl, heteroalkyl, heterocycyl, alkylaryl, alkylheterocyclic, or substituted derivatives thereof; and,

R_{10} is H, alkyl, heteroalkyl, heterocycyl, alkylaryl, alkylheterocyclic, or substituted derivatives thereof.

In an embodiment of the invention antimicrobial drugs that are selective inhibitors of peptide deformylase are used to treat infection while reducing or avoiding deleterious
10 side effects associated with therapeutic drugs that affect the host's biochemistry. In a preferred embodiment antimicrobial drugs of the invention are selective for peptide deformylase over other related enzymes such as angiotensin converting enzyme (ACE) thermolysin, collagenase and carboxypeptidase. Such selectivity minimizes the opportunity of adverse events associated with imbalances of these enzymes in the homostasis of tissue
15 and organ functions affected by these metalloproteases.

A preferred embodiment of the invention involves the use of compounds or compositions of the invention therapeutically or prophylactically against bacteria that are resistant to other antibiotics such as β -lactam, quinolone and vancomycin resistant bacteria.

In another embodiment, compounds of the invention may be used to treat
20 contaminated or infected items, such as crops, wood, metal or plastic and the like, by methods such as, but not limited to, spraying or dusting of that agent onto the contaminated item, or impregnating that agent into the item.


Additionally, the invention includes pharmaceutical compositions and therapies comprising a compound of the invention and a second antibacterial compound including
25 antibiotics of the following groups consisting of, but not limited to, aminoglycosides, amphenicols, ansamycins, beta-lactams, cephalosporins, cephamycins, monolactams, oxacephems, penicillins, lincosamides, macrolides, polypeptide antibiotics, tetracyclines, 2,4-diaminopyrimidines, nitrofurans, quinolones, streptogramins, sulfonamides, sulfones, oxazolidinones and glycyclines.

30


35

4. DETAILED DESCRIPTION OF THE INVENTION


The present invention relates to compounds represented by formulas I-IV:




Formula I



Formula II



Formula III



Formula IV

, or a pharmaceutically acceptable salt of such compounds, wherein:

Z is NHOH or OR_a wherein R_a is H, alkyl or a biocleavable moiety;

X is C=O or O=S=O;

Y is heteroalkyl, heterocyclic, or substituted derivatives thereof;

R₁ is C₁-C₁₀ alkyl, aryl, heterocyclic, heteroalkyl or a substituted derivative thereof;

R₂ and R₃ together with nitrogen represent a 4, 5, 6, or 7 membered heterocyclic ring optionally substituted with one of the following substituents -OH, -CH₂OH,

-O-C(=O)-heterocyclic, -O-C(=O)-NH-aryl, -NH-C(=O)-aryl,

-C(=O)-N(C₂-C₆ alkyl)₂, -NH-(C₁-C₆ alkyl)-heterocyclic,

-NH-C(=O)-NH-alkyl, -NH-C(=O)-NH-aryl, -NH-C(=O)-NH-heterocyclic;
or

R₂ and R₄ together with nitrogen form a ring through a -CH₂-CH₂- linkage wherein
R₃ is H, C₁-C₁₀ alkyl, aryl, heterocyclic, heteroalkyl or a substituted
5 derivative thereof; or R₂ is methyl;

R₃, as defined above forms a ring with R₂, or represents H, or is alkyl, heteroalkyl,
aryl, heterocyclic, and substituted derivatives thereof;

R₃ when not in a ring with R₂ is alkyl, heteroalkyl, aryl, heterocyclic, and substituted
derivatives thereof;

10 R₄ when not in a ring with R₂ is H, C₁-C₁₀ alkyl, aryl, heterocyclic, heteroalkyl or a
substituted derivative thereof; or is together with R₂ as defined above to form
a ring through a -CH₂-CH₂- linkage.

R₅ is -H, -NO₂, -NH₂, -NH-C(=O)-H, -NH-C(=O)-CH₃, -NH-S(O₂)-CH₃,
-CH₂-NH-alkyl, -CH₂-NH-heteroalkyl, CH₂-NH-heterocycyl, or substituted
15 derivatives thereof;

R₆ is -H, -NO₂, -NH₂, -NH-C(=O)-H, -NH-C(=O)-CH₃, -NH-S(O₂)-CH₃,
-CH₂-NH-alkyl, -CH₂-NH-heteroalkyl, -CH₂-NH-heterocycyl, or substituted
derivatives thereof;


20 R₇ or R₈ is -CHR₁₀-C(=O)-NH-OH; R₇ or R₈ when not -CHR-C(=O)-NH-OH is
alkyl, heteroalkyl, heterocyclic, alkylaryl, alkylheterocyclic, or substituted
derivatives thereof;

R₉ is H, alkyl, heteroalkyl, heterocycyl, alkylaryl, alkylheterocyclic, or substituted
derivatives thereof; and,

25 R₁₀ is H, alkyl, heteroalkyl, heterocycle, alkylaryl, alkylheterocyclic, or substituted
derivatives thereof.

A subset of preferred compounds encompassed by Formula I and II is defined
wherein R₂ and R₃ together with the nitrogen form a 4 membered azetidiny ring through a
-CH₂-CH₂-CH₂- linkage which is optionally substituted with one of the following
substituents: -OH, -CH₂OH, -O-C(=O)-heterocyclic, -O-C(=O)-NH-aryl, -NH-C(=O)-aryl,
30 -C(=O)-N(C₂-C₆ alkyl)₂, -NH-(C₁-C₆ alkyl)-heterocyclic, -NH-C(=O)-NH-alkyl,
-NH-C(=O)-NH-aryl, -NH-C(=O)-NH-heterocyclic. Compounds of this subset may be
represented Formula V.

5




Formula V

10

A subset of preferred compounds encompassed by Formulas I and II is defined wherein R_2 and R_3 together with the nitrogen form a five membered pyrrolidinyl ring through a $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ linkage or a five membered thioproline ring through a $-\text{CH}_2-\text{CH}_2-\text{S}-\text{CH}_2-$ linkage which is optionally substituted with one of the following substituents: OH, CH_2OH , $-\text{O}-\text{C}(=\text{O})$ -heterocyclic, $-\text{O}-\text{C}(=\text{O})$ -NH-aryl, $-\text{NH}-\text{C}(=\text{O})$ -aryl, $-\text{C}(=\text{O})$ -N(C_1 - C_6 alkyl) $_2$, $-\text{C}(=\text{O})$ -NH-(C_1 - C_6 alkyl), $-\text{C}(=\text{O})$ -heterocyclic, $-\text{NH}-(\text{C}_1$ - C_6 alkyl)-heterocyclic, $-\text{NH}-\text{C}(=\text{O})$ -NH-alkyl, $-\text{NH}-\text{C}(=\text{O})$ -NH-aryl, $-\text{NH}-\text{C}(=\text{O})$ -NH-heterocyclic. Compounds of this subset may be represented by Formulas VI, VII, VIII, and IX.


20



25


Formula VI

30




35

Formula VII




Formula VIII




Formula IX

A subset of preferred compounds encompassed by Formulas I and II is defined wherein R_2 and R_3 together with the nitrogen form a six membered piperidiny ring through a $-CH_2-CH_2-CH_2-CH_2-CH_2-$ linkage which is optionally substituted with one of the following substituents: $-OH$, $-CH_2OH$, $-O-C(=O)-$ heterocyclic, $-O-C(=O)-NH$ -aryl, $-NH-C(=O)-$ aryl, $-C(=O)-N(C_2-C_6 \text{ alkyl})_2$, $-NH-(C_1-C_6 \text{ alkyl})$ -heterocyclic. Compounds of this subset may be represented Formulas X, XI, XII, and XIII.



Formula X


5



Formula XI

10


15



Formula XII


20

25




Formula XIII

A subset of preferred compounds encompassed by Formulas I and II is defined wherein R_1 is selected from the group consisting of 3,4-difluorophenyl, phenyl and cyclohexyl; and, R_2 and R_3 together with the nitrogen form a six membered piperidiny ring through a $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ linkage which is optionally substituted with one of the following substituents: $-\text{OH}$, $-\text{CH}_2\text{OH}$, $-\text{O}-\text{C}(=\text{O})$ -heterocyclic, $-\text{O}-\text{C}(=\text{O})$ -NH-aryl, $-\text{NH}-\text{C}(=\text{O})$ -aryl, $-\text{C}(=\text{O})-\text{N}(\text{C}_2-\text{C}_6 \text{ alkyl})_2$, $-\text{NH}-(\text{C}_1-\text{C}_6 \text{ alkyl})$ -heterocyclic, $-\text{NH}-\text{C}(=\text{O})$ -NH-alkyl, $-\text{NH}-\text{C}(=\text{O})$ -NH-aryl, $-\text{NH}-\text{C}(=\text{O})$ -NH-heterocyclic. Compounds of this subset may be represented by Formula XIV.




Formula XIV

A subset of preferred compounds encompassed by Formula I is defined in Formula XV wherein R is selected from the group consisting of alkyl, heteroalkyl, aryl, heterocyclic, -NH-C(=O)-NH-alkyl, -NH-C(=O)-NH-aryl, -NH-C(=O)-NH-heterocyclic, and substituted derivatives thereof. Compounds of this set may be represented by Formula XV.




Formula XV

A subset of preferred compounds encompassed by Formula III is defined by Formulas XVI and XVII wherein R is selected from the group consisting of alkyl, heteroalkyl, aryl, heterocyclic, and substituted derivatives thereof. Compounds of this subset may be represented by Formulas XVI and XVII.



Formula XVI




5

Formula XVII

A subset of preferred compounds encompassed by Formula III is defined by Formula XVIII wherein R is selected from the group consisting of -H, -NO₂, -NH₂, -NH-(C=O)-CH₃, -NH-SO₂-CH₃, alkyl, and heteroalkyl. Compounds of this subset may be represented by

10 Formula XVIII.




15

Formula XVIII

A subset of preferred compounds encompassed by Formula IV is defined by

20 Formulas XIX and XX wherein R₇, R₈, R₉, and R₁₀ are selected from the group consisting of -H, -NO₂, -NH₂, -NH-(C=O)-CH₃, -NH-SO₂-CH₃, alkyl, heteroalkyl, aryl, heterocycyl, and alkylaryl. Compounds of this subset may be represented by Formula XIX.




25

Formula XIX

30


35



Formula XX


10

A subset of preferred compounds encompassed by Formula III is defined by Formula XXI, wherein M is 0 or 1, R is alkyl, the stereochemistry at the center indicated with * is R, or S, and Z is OH or NH-OH.



Formula XXI

20 A subset of preferred compounds encompassed by Formula IV is defined by Formula XXII, wherein R₉ is selected from the group consisting of H, alkyl, heteroalkyl, aryl, heterocycyl, and alkylaryl; R₁₀ is selected from the group consisting of H, alkyl, heteroalkyl, aryl, heterocycle, and alkylaryl; and R₁₁ is selected from the group consisting of H, alkyl, heteroalkyl, aryl, heterocycle, and alkylaryl.



Formula XXII

30

Definitions

Throughout the instant application the following terms are defined below unless otherwise stated.

35

As used herein the terms "alkyl" means a straight or branched chain alkyl moiety including, but not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, and isohexyl.

As used herein the term "alkenyl" means a straight or branched chain hydrocarbon moiety having one or more double bonds including, but not limited to, propene, 1-butene, 2-butene, iso-butene, 1-pentene, 2-pentene, iso-pentene, 1-hexene, 2-hexene, and 3-hexene.

As used herein the term "cycloalkyl" means a saturated alicyclic moiety having one or more fused ring systems with from 3-12 carbons including, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

As used herein the term "heteroalkyl" means an optionally substituted alkyl chain containing one or more heteroatoms or their oxidized forms selected from nitrogen, sulfur, oxygen, or phosphorous.

As used herein the terms "heterocycle" or "heterocyclic", means a 5 to 7 member ring, either aromatic or non-aromatic, containing one or more heteroatoms selected from nitrogen, sulfur, and oxygen, and optionally substituted or fused to a benzene ring system examples, include but are not limited to, thiazolyl, imidazolyl, oxazolyl, indoyl, morpholinyl, pyridinyl, pyrrolidinyl, pyrimidinyl, thienyl, furyl, piperidinyl, and piperazinyl.

As used herein the term "aryl" refers to aromatic rings and substituted derivatives thereof including, but not limited to, phenyl and fused ring systems like naphthyl.

As used herein the terms "biolabile" or "biocleavable" moiety refers to a pharmaceutically acceptable, biologically degradable moiety or derivative of a compound of the invention, that is a prodrug which, upon administration to an animal or human being, is converted in the body to a compound of the invention. Such biolabile or biocleavable moieties are particularly advantageous in providing compounds of the invention that are suitable for oral administration. The suitability of any particular biolabile or biocleavable group can be assessed by conventional *in vivo* animal or *in vitro* enzyme hydrolysis studies.

It should be recognized that there are several actual or potential chiral centers in the compounds according to the invention due to the presence of asymmetric carbons. A chiral carbon within a compound gives rise to stereoisomers with R or S stereochemistry at each chiral center. The invention includes all such isomers, stereoisomers, diastereomers, enantiomers, and all racemic or optically pure forms of the compounds.


One of ordinary skill in the art of organic or medicinal chemistry will know how to resolve racemic compounds or to otherwise purify a desired stereoisomer using chiral techniques or separations.

Because of possible discrepancies in using chemical nomenclature where structures are provided for compounds or moieties the structure controls the definition of the compound or moiety, and not the chemical name.



Representative examples of the compounds of formula I are shown below in Tables 1-13, these examples are not meant to be limiting but are for illustrative purposes: the IC_{50} data is the concentration (nanomolar) of the compound that inhibits E coli peptide deformylase at 50%, and the MIC data refers to the concentration of compound in micrograms per milliliters to achieve an inhibition of growth of a test organism such as E. coli.








Table 1







Pyrrolidine Carbamate Series I









Formula VI







Cmpd. No.	R	R'	$IC_{50}(nM)$	MIC ($\mu g/ml$)
1	 2-Chlorophenyl	H	17	12.5
2	 3-Chlorophenyl	H	14	50






Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
3	 4-Chlorophenyl	H	16	50
4	 2-Fluorophenyl	H	13	25
5	 3-Fluorophenyl	H	21	25
6	 4-Fluorophenyl	H	21	12.5
7	 2-Bromophenyl	H	32	25
8	 3-Bromophenyl	H	22	25
9	 Phenyl	H	19	25








Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
10	 4-Methoxyphenyl	H	9	25
11	 4-Trifluoromethoxyphenyl	H	40	50
12	 4- (N,N-Dimethyl-amino)phenyl	H	21	25
13	 4-Bromophenyl	H	33	50
14	 4-Methylphenyl	H	31	25
15	 4-n-Butylphenyl	H	160	25



Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
16	 4-Trifluoromethylphenyl	H	25	>100
17	 2-Methoxyphenyl	H	27	25
18	 2-Ethylphenyl	H	28	12.5
19	 2-i-Propylphenyl	H	32	12.5
20	 3-Methylphenyl	H	17	12.5
21	 3-Nitrophenyl	H	25	25

35

Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
22	 3-Methylmercaptophenyl	H	27	25
23	 4-Chloro-3-nitrophenyl	H	37	12.5
24	 4-Chloro-3-trifluoromethyl- phenyl	H	103	25
25	 2,4-Dichlorophenyl	H	65	25
26	 2,4-Dimethoxyphenyl	H	16	12.5
27	 4-Chloro-2-trifluoromethyl- phenyl	H	54	50


Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
28	 2-5-Difluorophenyl	H	10	25
29	 2-Methoxy-5-chlorophenyl	H	38	25
30	 2-Methoxy-5-methylphenyl	H	26	25
31	 3,5-Dimethyl-phenyl	H	129	100
32	 3,5-bis-(Trifluoromethyl)phenyl	H	350	50

Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
33	 3,5-Dichlorophenyl	H	226	50
34	 4-(2,6-Dichloropyridyl)	H	26	100
35	 <i>n</i> -Propyl	H	19	12.5
36	 Benzyl	H	5	----
37	 2-(Ethyl-2-thiophene)	H	29	12.5
38	 4-Phenoxyphenyl	H	39	25
39	 4-Trifluoromethanemercapto- phenyl	H	37	25

Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
40	 2-Phenylphenyl	H	43	25
41	 2-Phenoxyphenyl	H	54	25



*Although the stereochemistry at the 3-position of the pyrrolidine ring is not specified, one of ordinary skill in the art would readily recognize that the stereocenter may be generated in racemic form, *i.e.*, a mixture of *R* and *S* enantiomers or with the desired stereochemical orientation, *i.e.*, the substantially or isolated *R* or *S* enantiomer.

Table 2
Pyrrolidine Carbamate Series II




Formula VII

Cmpd. No.	R	IC ₅₀ (nM)	MIC (μg/ml)
42	 N-Pyrrolidine	11	6
43	 N-Pyrrolidine-2 (R) -methanol	14	50
44	 N-Pyrrolidine-2 (S) -methanol	5	100
45	 N-Piperidine	8	6.25
46	 N-Homo-Piperidine	11	3.13






Cmpd. No.	R	IC ₅₀ (nM)	MIC (μg/ml)
47	 <i>N</i> -Morpholine	15	-----
48	 <i>N</i> -3 <i>S</i> -Pyrrolidinol	30	100




*Although the stereochemistry at the 3-position of the pyrrolidine ring is not specified, one of ordinary skill in the art would readily recognize that the stereocenter may be generated in racemic form, *i.e.*, a mixture of *R* and *S* enantiomers or with the desired stereochemical orientation, *i.e.*, the substantially or isolated *R* or *S* enantiomer.

Table 3
Piperidine Carbamate Series




Formula X

Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
49	 - Phenyl	H	90	50
50	 3,5-Dichlorophenyl	H	200	25
51	 3-Chlorophenyl	H	176	50
52	 4-Chlorophenyl	H	147	50
53	 4-Methoxyphenyl	H	51	25






Cmpd. No.	R	R'	IC ₅₀ (nM)	MIC (μg/ml)
54	 4-(N,N-Dimethylamino)phenyl	H	103	25
55	 4-Trifluoromethoxyphenyl	H	78	50
56	 4-Fluorophenyl	H	350	25









*Although the stereochemistry at the 3-position of the piperidine ring is not specified, one of ordinary skill in the art would readily recognize that the stereocenter may be generated in racemic form, *i.e.*, a mixture of *R* and *S* enantiomers or with the desired stereochemical orientation, *i.e.*, the substantially or isolated *R* or *S* enantiomer.



Table 4
Thioproline Series



Formula VIII


Cmpd. No.	R	IC ₅₀ (nM)	MIC (μg/ml)
57	 Isobutylamino	20	3.13
58	 Cyclopentylamino	12	1.56
59	 Cyclohexylamino	15	1.56
60	 Tetrahydrofurfurylamino	14	3.13
61	 Furfurylamino	17	1.56

Cmpd. No.	R	IC ₅₀ (nM)	MIC (μg/ml)
62		13	1.56
63		26	1.56
64		7	0.78
65		26	3.13
66		37	1.56
67		37	1.56
68		47	6.25
69		24	3.13

	Cmpd. No.	R	IC₅₀ (nM)	MIC (μg/ml)
5	70	 2,6-Dimethylmorpholino	21	6.25
10	71	 Thiomorpholino	25	3.13

*Although the stereochemistry at the 4-position of the thioproline ring is specified as the *S* enantiomer, one of ordinary skill in the art would readily recognize that the stereocenter may be generated in racemic form, *i.e.*, a mixture of *R* and *S* enantiomers, or with the *R* stereochemical orientation.


Table 5
Tic and Pyrrolidine Series



15

Cmpd. No.	R	IC ₅₀ (nM)
72	<p>20</p> <p>N-1,2,3,4-Tetrahydronathyl-1-amine 2-pyrrolidinamide</p>	22
73	<p>25</p> <p>N-1,2,3,4-Tetrahydronathyl-1-amine 2-pyrrolidinamide</p>	31
74	<p>30</p> <p>N-Pyrrolidine 2-pyrrolidinamide</p>	82, MIC=3.13 (μg/ml)

35

Cmpd. No.	R	IC ₅₀ (nM)
75	 Pyrrolidine 2-pyrrolidinamide	960

5

10

15


20

25

30

35

Table 6
Benzyloxypyrrolidine Series



Formula IX

Cmpd No.	R	IC ₅₀ (nM)
76	 2-Thiazolylamino	8
77	 Dimethylamino	57
78	 N-2-Methylpyridyl	40
79	 N-Cyclopentylamine	20
80	 N-Thiomorpholine	65

*Although the stereochemistry at the 2-position of the piperadine ring is specified, one of ordinary skill in the art would readily recognize that the stereocenter may be generated in racemic form, *i.e.*, a mixture of *R* and *S* enantiomers or with the desired stereochemical orientation, *i.e.*, the substantially or isolated *R* or *S* enantiomer.

5

10

15


20

25

30









35

Table 7
Homoproline Series I







Formula X

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
81	 Isobutylamino	13	12.5
82	 Cyclopentylamino	13	12.5
83	 Cyclohexylamino	13	12.5
84	 Tetrahydrofurfurylamino	11	6.25
85	 Furfurylamino	9	6.25

5	86		15	6.25
		2-Thienylmethylamino		
	87		18	6.25
		N-methyl-3-pyridyl		
10	88		20	6.25
		Benzylamino		
15	89		3	0.78
		2-Thiazolylamino		
20	90		17	1.56
		Dimethylamino		
25	91		15	1.56
		Piperidino		
	92		16	3.13
30		Homopiperidino		
	93		19	3.13
		Morpholino		

35

5	94		19	3.13
		2,6-Dimethylmorpholino		
10	95		17	1.56
		Thiomorpholino		
	96		19	6.25
		N-Methyl-N-piperazine		
15	97		13	1.56
		1-Pyrrolidinyl		


20 *Although the stereochemistry at the 2-position of the piperidine ring specified is as the S enantiomer, one of ordinary skill in the art would readily recognize that the stereocenter may be generated in racemic form, *i.e.*, a mixture of *R* and *S* enantiomers or as the *R* enantiomer.

25











30



35

Table 8
Homoproline II series



Formula XIV

Cmpd No.	R ₁	R ₂	IC ₅₀ (nM)
98	 3,4-Difluorophenyl	 1-Pyrrolidinyl	116
99	 Phenyl	 1-Pyrrolidinyl	22
100	 Cyclohexyl	 1-Pyrrolidinyl	9
101	 3,4-Difluorophenyl	 2-Thiazolylamino	81
102	 Phenyl	 2-Thiazolylamino	25

103			2
	Cyclohexyl	2-Thiazolylamino	.

5

*Although the stereochemistry at the 2-position of the piperidine ring is specified, one of ordinary skill in the art would readily recognize that the stereocenter may be generated in racemic form, *i.e.*, a mixture of *R* and *S* enantiomers or with the desired stereochemical orientation, *i.e.*, the substantially or isolated *R* or *S* enantiomer.

10

15


20

25






30









35






Table 9
Nipecotic Acid Series



Formula XII


Cmpd. No.	R	IC ₅₀ (nM)	MIC (μg/ml)
104	 Isobutylamino	96	50
105	 Cyclopentylamino	62	50
106	 Cyclohexylamino	38	25
107	 Tetrahydrofurfurylamino	41	-----
108	 Furfurylamino	150	50

Cmpd. No.	R	IC ₅₀ (nM)	MIC (μg/ml)
109	 Thienylmethylamino	30	6.25
110	 N-2-Methylpyridyl	26	-----
111	 Benzylamino	25	6.25
112	 2-Thiazolylamino	116	100
113	 Dimethylamino	31	50
114	 Pyrrolidinyl	40	-----
115	 Piperidino	112	50
116	 Homopiperidino	44	100

Cmpd. No.	R	IC ₅₀ (nM)	MIC (μg/ml)
117	 Morpholino	45	100
118	 2,6-Dimethylmorpholino	49	100
119	 Thiomorpholino	205	6.25
120	 N-Methyl-N-piperazine	67	-----
120A	 Diethylamino	50	-----









*Although the stereochemistry at the 3-position of the piperidine ring is not specified, one of ordinary skill in the art would readily recognize that the stereocenter may be generated in racemic form, *i.e.*, a mixture of *R* and *S* enantiomers or with the desired stereochemical orientation, *i.e.*, the substantially or isolated *R* or *S* enantiomer.


Table 10
Isonipectic Acid Series



Formula XIII

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
123	 Isobutylamino	48	50
124	 Cyclopentylamino	81	25
125	 Furfurylamino	44	100
126	 2-Thienylmethylamino	129	25
127	 N-2-methylpyridyl	34	50

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
128	 N-Benzyl	48	50
129	 2-Thiazolylamino	29	-----
130	 Dimethylamino	109	-----
131	 1-Pyrrolidinyl	135	-----
132	 Piperidino	318	100
133	 Homopiperidino	85	-----
134	 Morpholino	83	-----
135	 2,6-Dimethylmorpholino	74	-----

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
136	 Thiomorpholino	72	-----

5

10

15


20

25

30






35





Table 11
Related Peptide Deformylase Inhibitors












Formula XV






Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
137	 Valinyl-pyrrolidinamide	6.1	6.25
138	 Amino-valeryl-N,N-dimethylamide	7.7	-----
139	 Phenylalanyl-N,N-dimethylamide	10.1	25








Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
140		5.5	-----
141		5.5	50
142		10.1	50
143		5.4	12.5
144		11.4	25







Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
145		6.0	50
146		6.4	25
147		15	25
148		1450	-----






Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
149		64.0	-----
	Amino-valeryl-piperidinamide		
150		114.7	100
	Phenylalanyl-piperidinamide		
151		222.6	-----
	Methionyl-piperidinamide		
152		101.5	25
	O-Benzyl-serinyl-piperidinamide		





Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
153	 O-Benzyl-tyrosinyl-piperidinamide	96.5	-----
154	 Valinyl-benzylamide	34.1	25
155	 Amino-Valeryl-benzylamide	7.4	25
156	 Phenylalanyl-benzylamide	6.8	-----
157	 Methionyl-benzylamide	26.7	-----





Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
158	 O-Benzyl-Tyrosinyl-benzylamide	6.0	-----
159	 O-Benzyl-Tyrosinyl-benzylamide	8.6	-----
160	 Phenylalanyl-anilnamide	73	100
161	 N-Butylamine	920.5	100
162	 N-Pentylamine	159.8	100







Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
163		523.4	100
164		179	100
165		120.6	100
166		143	100
167		408.8	100
168		1423	100
169		1119	100







	Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
5	170	 N-4-Fluoroaniline	457.7	100
10	171	 N-3-Bromoaniline	576	100
15	172	 N-4-Bromoaniline	577	100
20	173	 N-4-Chloroaniline	542	-----
25	174	 N-3,5-Dimethoxyaniline	547	-----
30	175	 N-3-Phenoxyaniline	1393	-----







Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
176	 N-4-Phenoxyaniline	669	100
177	 N-(4-N-Acetamidyl)-aniline	2553	100
178	 Serinyl-Dimethyl amide	10.9	100
179	 Lysinyl-pyrrolidinamide	7.1	100
180	 Glycinyl-(2-acetohydroxamate)-pyrrolidinamide	13.2	100








Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
181		39.9	100
182		8.7	100
183		5.1	-----
184		146	100

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
185		35.9	100
	Lysinyl-benzyl amide		
186		52	100
	Glycyl-4-methoxy-anilnamide		
187		11	---
	Lysinyl-4-methoxy-anilnamide		
188		35	100
	Glycyl (2-acetohydroxyamate) 4-methoxyanilnamide		

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
189	 <p>Glycyl-(2-propiohydroxyamate) 4-methoxyanilamide</p>	8	100
190	 <p>Glycyl-(4-fluoro)-anilamide</p>	42	-----
191	 <p>N-Propionate (4-fluoro)-anilamide</p>	172	100
192	 <p>N-2-Piperidine-methanol</p>	30	-----
193	 <p>N-4-Piperidine-ethanol</p>	17	-----
194	 <p>N-Benzylethanolamine</p>	14	-----

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
5 195	 N,N-Diethanolamine	200	-----
10 196	 N-Methyl-N-[(3,4-Dimethoxy)-2-phenylethyl]-amine	58	-----
15 197	 N-2-(Methylamino)ethanol	41	-----
20 198	 N-3-Aminobenzyl alcohol	215	-----
25 199	 N-4-Aminophenylethyl alcohol	122	-----
30 35 200	 N-(N-Benzyl glyciny)-2-ethanoic acid amine	2030	-----

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
201	 N-2-Piperidine-methanol	30	-----
202	 N-4-Piperidine-ethanol	17	-----
203	 N-Benzyl-ethanolamine	14	-----
204	 N-Diethanolamine	200	-----
210	 N-[N-Ethyl-(2-(3,5-dimethoxyphenyl))-N-methyl]amine	58	-----
211	 N-2-(Methylamino) ethanol	41	-----

Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
212	 N-3-Aminobenzyl alcohol	215	-----
213	 N-4-Aminophenylethyl alcohol	122	-----
214	 3-Pyrrolidinol	89	50
215	 2-Pyrrolidinol	80	25
216	 3-Piperidinol	35	50
217	 N-Piperazine-N-benzylamide	58	100
218	 N-Piperazine-N-pyrrolidinamide	47	50













Cmpd No.	R	IC ₅₀ (nM)	MIC (μg/ml)
219	 N-Pyrrolidine-2(S)-benzylamide	23	100
220	 N-Pyrrolidine-3(R)-benzylamide	39	50
221	 N-Azetidine-(3-O-anilnamide)	33	50
222	 N-3-Benzylamide-azetidine	17	50
223	 N-Piperidine-2-pyrrolidinamide	194	100

Table 12
Capped Valine Series I

5


**Formula XVI**

10

Cmpd No.	R	IC ₅₀ (nM)
224		3000
225	 -Butyl	5000
226	 Ethyl 2-phenyl	4500
227	 Cyclopentyl	3500
228	 2-Methyltetrahydrofuran	3500
229	 4-Chlorophenyl	5800







35

Table 13
Capped Valine Series II




Formula XVII

10

Cmpd No.	R	IC ₅₀ (nM)
230	 1-Naphthalene-1,2,3,4-tetrahydride	2300
231	 n-Butyl	5200
232	 Ethyl 2-phenyl	4000
233	 Cyclopentyl	1900
234	 2-Methyltetrahydrofuran	1800
235	 4-Chlorophenyl	2000

15
20
25
30
35


Table 14
Tetrahydrobenzothiophene Series



Formula XVIII

Cmpd. No.	R _A	R _B	IC ₅₀ (nM)
239	-NO ₂	H	13000
240	H	-NO ₂	10000
241	H	-NHC(=O)CH ₃	6000
242	H	-NHC(=O)H	6000
243	H	-NHS(O ₂)CH ₃	9000


Table 15
Hydantoin Series I



Formula XIX

Cmpd. No.	R ₉	R ₈	R ₁₀	IC ₅₀ (nM)
244	-CH ₂ CH(CH ₃) ₂	H	-(CH ₂) ₂ S(CH ₃)	59,000
245	-(CH ₂) ₂ S(CH ₃)	H	-(CH ₂) ₂ S(CH ₃)	127000
246	-CH(CH ₃) ₂	H	-(CH ₂) ₂ S(CH ₃)	272000
300	-CH ₂ (C ₆ H ₅)	H	-(CH ₂) ₂ S(CH ₃)	13000
247	-CH ₂ (C ₆ H ₅)	H	-CH ₂ (C ₆ H ₅)	19000
248	-(CH ₂) ₃ CH ₃	H	-(CH ₂) ₂ S(CH ₃)	51000
249	-(CH ₂) ₃ CH ₃	H	-CH ₂ C ₆ H ₅	94000
251	-CH(CH ₃) ₂	H	-CH(CH ₃) ₂	185000
252	-(CH ₂) ₂ S(CH ₃)	H	-CH ₂ CH(CH ₃) ₂	74000
253	-CH ₂ (C ₆ H ₅)	H	-CH ₂ CH(CH ₃) ₂	32000
254	-(CH ₂) ₃ CH ₃	H	-CH ₂ CH(CH ₃) ₂	60000


Table 16
Hydantoin Series II



Formula XX

Cmpd. No.	R ₇	R ₉	R ₁₀	IC ₅₀ (nM)
255	-CH ₃	-CH(CH ₃) ₂	H	133000
256	-CH ₂ (C ₆ H ₅)	-(CH ₂) ₃ (CH ₃)	H	3000
257	-CH ₃	-(CH ₂) ₂ S(CH ₃)	H	47000
258	-CH ₃	-CH ₂ CH(CH ₃) ₂	H	32000
259	-CH ₃	-CH ₂ (C ₆ H ₅)	H	76000
260	-CH ₃	-(CH ₂) ₃ CH ₃	H	24000
250	-CH ₂ C(=O)NH-OH	-CH(CH ₃) ₂	H	75000
261	-CH ₂ C ₆ H ₅	-CH(CH ₃) ₂	H	1000
262	-CH ₂ C ₆ H ₅	-CH ₂ CH(CH ₃) ₂	H	2000
263	-CH ₂ C ₆ H ₅	-CH ₂ C ₆ H ₅	H	88000
264		-CH ₂ CH(CH ₃) ₂	H	6520
264A		-CH ₂ CH(CH ₃) ₂	H	5000


Table 17
Hydantoin Series III



Formula XXII





Cmpd No.	R ₉	R ₁₀	R ₁₁	IC ₅₀ (nM)
266	-CH ₂ CH(CH ₃) ₂	H	N-Pyrrolidine	39000
267	-CH ₂ CH(CH ₃) ₂	H	N-Homopiperidine	10000
301	-CH ₂ CH(CH ₃) ₂	-CH(CH ₃) ₂	N-Pyrrolidine	1000
268	-CH ₂ CH(CH ₃) ₂	-CH(CH ₃) ₂	N-Homopiperidine	2000
269	-(CH ₂) ₃ CH ₃	-CH(CH ₃) ₂	N-Homopiperidine	1000
270	-CH ₂ CH(CH ₃) ₂	H	N-2-Pyrrolidine-methanol	35000
271	-CH ₂ CH(CH ₃) ₂	-CH(CH ₃) ₂	N-2-Pyrrolidine-methanol	190
272	-CH ₂ CH(CH ₃) ₂	-CH ₂ (C ₆ H ₅)	N-2-Pyrrolidine-methanol	400
273	-CH ₂ CH(CH ₃) ₂	-CH ₂ (C ₆ H ₅)	N-2-Pyrrolidine-methanol	800

Table 18
Capped Dipeptide Series III






Cmpd No.	R	0	M	Z	IC ₅₀ (nM)
274	-(CH ₂) ₃ CH ₃	L	0	-NH-OH	42
275	-(CH ₂) ₃ CH ₃	D	0	-NH-OH	502
276	-CH ₂ (C ₆ H ₁₁)	L	0	-NH-OH	110
277	-(CH ₂) ₃ CH ₃	L	1	-NH-OH	100
278	-(CH ₂) ₃ CH ₃	D	1	-NH-OH	1700
279	-(CH ₂) ₃ CH ₃	D	1	-OH	1600

Table 19
Related Deformylase Inhibitors

	Cmpd No.	Structure	IC₅₀ (nM)	MIC (μg/ml)
5	310		103000	-----
10				
15	311		180	50
20	312		47	25
25				
30	313		10	12.5

35

5	314		40	12.5
10	315		30	6.25
15	316		60	25
20				

25

30

35

4.9 Biological Assays

The compounds of the invention have utility as agents effective against a variety of species of bacteria, including infectious pathogenic bacteria. The invention also includes novel pharmaceutical compositions which comprise the compounds of the invention
5 formulated in pharmaceutically acceptable formulations.

One embodiment the invention features a method for treating a subject infected with an infectious agent by administering to that subject a therapeutically effective amount of a compound of the invention which causes PDF inhibition in the infectious agent as determined by the assays of the invention. Veterinary uses are also included in this
10 embodiment, as described below. Such administration can be by any method known to those skilled in the art, for example, by topical application or by systemic administration. Additional antibacterial compounds may be adjunctively administered, as described below.

In yet another embodiment, compounds of the invention can be used to treat contaminated or infected items, such as crops, wood, metal or plastic and the like, by
15 methods such as, but not limited to, spraying or dusting of that agent onto the contaminated item, or impregnating that agent into the item.

By "therapeutically effective amount" or "prophylactically effective" is meant an amount that relieves (to some extent) one or more symptoms of the disease or condition in the subject. Additionally, by "therapeutically effective amount" is meant an amount that
20 returns to normal, either partially or completely, physiological or biochemical parameters associated with or causative of a bacterial disease or condition.

4.10 Selectivity Profile Assays

High through-put screens described in U.S. application number 09/449,419 filed
25 November 29, 1999, which is incorporated by reference in its entirety, are used to determine the binding affinity of compounds to PDF in a novel assay that utilizes the enzyme's native iron metal center and the binding affinity to several other metalloproteases including thermolysin, carboxypeptidase, collagenase and angiotensin converting enzyme. The selectivity profile is then determined by calculating the ratio of the binding affinity of the
30 inhibitor to: (1) peptide deformylase with its native iron metal center and (2) other metalloproteases. It is preferred that compounds bind to PDF with a greater than or equal to 10-fold affinity relative to the other metalloproteases. It is highly preferred that compounds bind to PDF with a greater than or equal to 100-fold affinity relative to the other metalloproteases. In another embodiment, it is preferred that compounds bind to PDF with
35

a greater than or equal to 100-fold affinity relative to thermolysin and greater than or equal to 300-fold relative to carboxypeptidase, collagenase and angiotensin converting enzyme.

The screening methods of the invention are a substantial improvement over prior technologies in that: (1) the inhibition of peptide deformylase is determined using the
5 enzyme's native iron catalytic metal center and (2) the risk of deleterious side effects can be minimized since the novel inhibitors identified by the means of the invention interact negligibly with important metalloproteases of the host's biochemical machinery. The metalloproteases that are tested include thermolysin, carboxypeptidase, collagenase and angiotensin converting enzyme.

10 Initial high through-put assays are performed to determine the percent Fe-PDF inhibition at a particular concentration of test compound. The concentration of test compound is typical between 1 μ M - 5 mM. While the concentration may be varied, a preferred concentration of test compounds is 300 μ M. The concentration of reagents described below reflects assays which contain 300 μ M of test compound. However, the
15 concentration of reagents can be easily adjusted to accommodate the concentration of test compound given the parameters provided by the invention in the disclosure below. The reactions are set up in polypropylene plates. Preferred plates have 96- or 384-wells. A preferred plate is a 96-well polypropylene plate with 12 columns and 8 rows. One column is reserved for the negative control (20% DMSO), one column is for the positive control
20 (actinonin, Sigma, St. Louis, MO) and the rest are for test compounds. The reaction mixture, containing CHELEXTM 100 (Sigma, St. Louis, MO) treated water, 10.6 mM NaCl, 1.06 mg/ml BSA, 10.6 mM TCEP-HCl (Pierce, Rockford IL.) 35.3 mM fMAS, 23.5 mM NAD and 17.6 U/ml formate dehydrogenase (FDH) (Roche Molecular Biochemicals, Indianapolis, IN) is agitated and approximately 85 μ l of the reaction mixture is then added
25 to each well on the reaction plate. Endpoint data is recorded in a spectrophotometer by monitoring at 335-345 nanometers. 340 nanometers is preferred. To initiate the reactions 10 μ l of 1/20,000 fold diluted Fe-PDF (Rajagopalan et al., (1997) *J. Am. Chem. Soc.* 119:12418-12419; Rajagopalan et al., (1997) *Biochemistry* 36:13910-13918, both of which are incorporated by reference) is added to each well. The TCEP buffer is necessary for
30 stabilizing the diluted enzyme. The reaction plate is then mixed on a vortex mixer at low speed. Next the reactions are monitored in a spectrophotometer. A preferred spectrophotometer is the Spectromax 250 (Molecular Devices Corp., Sunnyvale, CA). Analysis of results from these experiments allows the determination of percent inhibition of Fe-PDF at 300 μ M test compound.

35

The assays to determine the binding affinity of the test compounds are based on the results obtained from the percent Fe-PDF inhibition studies. The initial concentrations of the test compounds in the binding affinity assays are adjusted according to the results in the percent Fe-PDF inhibition studies.

5 Compounds are initially prepared in a 96-well working plate with 12 columns and 8 rows. Again, the invention is not limited to this size plate, one skilled in the art can choose plate size appropriate to ones needs. One column is for the negative control (20% DMSO), another is reserved for the serial dilutions of the positive control, actinonin, and the rest are for the test compounds. The test compounds are diluted from 1/5 to 1/640 fold. 85 μ L of
10 reaction mixture is added to each well along with 5 μ L of the appropriately diluted test compound. The reactions are initiated by addition of 10 μ L of 1/20,000 fold dilute Fe-PDF. After addition of the Fe-PDF the plate is immediately inserted into the spectrophotometer and monitored at 335-345 nanometers and preferably 340 nanometers. Analysis of the data provided from the spectrophotometer readings gives the binding affinities for the test
15 compounds.

The thermolysin assay is designed to detect compounds that have an inhibitory effect on the metalloprotease thermolysin. The assay is based on the use of succinylated casein in conjunction with TNBSA (trinitrobenzene sulfonic acid)(Hatakeyama et al., (1992) *Anal. Biochem.* 204:181-184; Bubnis et al., (1992) *Anal. Biochem.* 207:129-133; Habeeb et al.,
20 (1996) *Anal. Biochem.* 14:328-336). Native casein has been treated with succinic anhydride to block available primary amines on the surface of the protein. Thermolysin acts to cleave the peptide bonds to expose primary amines. TNBSA reacts with primary amines to produce a yellow-orange color that can be detected at A430 and quantitated. A control well consisting of substrate, enzyme, and TNBSA with 1% DMSO is used to determine the basal
25 signal level. Any test well with a relative signal lower than the control level would indicate that thermolysin activity has been disrupted. The known inhibitor of thermolysin, phosphoramidon, is added for the calculation IC_{50} data (*i.e.*, the concentration of test compound that results in a fifty percent reduction of enzyme activity).

The carboxypeptidase A assay is designed to detect compounds that have an
30 inhibitory effect on carboxypeptidase A. The assay is based on the use of a blue shift of the absorption spectrum that occurs upon the hydrolysis of a furanacryloyl peptide FAPP (FA-Phe-Phe-OH) by the enzyme (Peterson et al., (1982) *Anal. Biochem.* 125:420-426, which is incorporated by reference in its entirety). Activity by the enzyme will produce a decrease in OD at A330 between an initial reading at time zero and a second reading after
35 one hour. A control-well consisting of enzyme, substrate, and 1% DMSO is used to

determine a maximum OD. Any test well with an OD lower than the control well will indicate that the enzyme activity has been disrupted. A known inhibitor of Carboxypeptidase A from potato tubers is added for the calculation of IC₅₀ data.

The collagenase assay is designed to detect compounds that have an inhibitory effect
5 on bacterial collagenase. The assay is based on the use of a blue shift of the absorption spectrum that occurs upon the hydrolysis of a furanacryloyl peptide FALGPA (FA-Leu-Gly-Pro-Ala-OH) by the enzyme (Van Wart et al., (1981) *Anal. Biochem.* 113:356-365, which is incorporated by reference in its entirety). Activity by the enzyme will produce a decrease in OD at A330 between an initial reading at time zero and a second
10 reading after twenty minutes. A control-well consisting of enzyme, substrate, and 1% DMSO is used to determine a maximum OD. Any test well with a OD lower than the control well would indicate that the enzyme activity has been disrupted. A peptide that is a known inhibitor of collagenase is used for calculation if IC₅₀'s.

The angiotensin converting enzyme (ACE) assay is designed to detect compounds
15 that have an inhibitory effect on the enzyme ACE. Angiotensin converting enzyme is a halide-activated peptidase that splits off hydrolytically the dipeptide His-Leu from the carboxyl end of the decapeptide angiotensin I (Asp-Arg-Val-Try-Ile-His-Pro-Phe-His-Leu), thereby converting it to the vasopressor octapeptide angiotensin II. The assay utilizes the internally quenched fluorescent tripeptide derivative Abz-Gly-Phe(NO₂)-Pro. The
20 fluorescent aminobenzoyl (Abz) group incorporated in this tripeptide can be excited in the 300-380 nm range upon cleavage of the peptide bond Gly-Phe(NO₂)-Pro (Carmel and Yaron, (1978) *Anal. Biochem.* 87:265-273, which is incorporated by reference in its entirety). Fluorescence is proportional to the amount of liberated Abz-Gly, and a percent control value can be determined by comparing the fluorescence of a sample compound well
25 to that of a control well containing only 1% DMSO. The substrate has a published $K_m = 0.21 \text{ mM} \pm 0.1 \text{ mM}$.

During the course of screens designed to identify inhibitors of peptide deformylase, it was discovered by the present inventors that a variety of compounds inhibit peptide deformylase with a greater than or equal to 10-fold affinity, and preferably a greater than or
30 equal to 100-fold affinity, relative to other metalloproteases.

Determination of MIC

The minimum inhibitory concentration (MIC) against bacterial organisms was determined certain compounds noted herein. Methods known in the art may be used such as
35 broth microdilution testing, using a range of concentrations of each test compound (1993,

National Committee for Clinical Laboratory Standards, Methods for Dilution Antimicrobial Susceptibility Tests For Bacteria That Grow Aerobically - Third Edition: Approved Standard, M7-A3, which is incorporated by reference herein in its entirety). The MIC against a variety of pathogens are determined using the same method. Pathogenic species to be tested generally include, but are not limited to: *E. coli*, *Enterococcus faecium*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Staphylococcus epidermis*, *Shigella flexneri*, and *Salmonella typhimurium*.

10 Formulations

The compounds described above can be provided as pharmaceutically acceptable formulations using formulation methods known to those of ordinary skill in the art. Administration as used in the invention includes those suitable for oral, ophthalmic, (including intravitreal or intracameral), topical, mucosal (including buccal, rectal, vaginal, nasal and sublingual), transdermal or parenteral (including subcutaneous, intramuscular, intravenous, bolus injection, intradermal, intratracheal, by implantable pump, and epidural) administration. In addition, the combinations may be incorporated into biodegradable polymers allowing for sustained release of the compound, the polymers being implanted in the vicinity of where drug delivery is desired. Biodegradable polymers and their use are described, for example, in detail in Brem et al., *J. Neurosurg.* 74:441-446 (1991).

The formulations include those suitable for oral, rectal, ophthalmic, (including intravitreal or intracameral) nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intratracheal, and epidural) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil emulsion and as a bolus, etc.

Formulations suitable for topical administration in the mouth include lozenges comprising the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the ingredient to be administered in a
5 suitable liquid carrier.

Formulations suitable for topical administration to the skin may be presented as ointments, creams, gels and pastes comprising the ingredient to be administered in a pharmaceutical acceptable carrier. A preferred topical delivery system is a transdermal patch containing the ingredient to be administered.

10 Formulations for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or a salicylate.

Formulations suitable for nasal administration, wherein the carrier is a solid, include a coarse powder having a particle size, for example, in the range of 20 to 500 microns which is administered in the manner in which snuff is administered, *i.e.*, by rapid inhalation
15 through the nasal passage from a container of the powder held close up to the nose.

Suitable formulations, wherein the carrier is a liquid, for administration, as for example, a nasal spray or as nasal drops, include aqueous or oily solutions of the active ingredient.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the
20 active ingredient such carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include
25 suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) conditions requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the
30 kind previously described.

Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the administered ingredient.

It should be understood that in addition to the ingredients, particularly mentioned
35 above, the formulations of the present invention may include other agents conventional in

the art having regard to the type of formulation in question, for example, those suitable for oral administration may include flavoring agents.

Oral dosage forms include tablets, capsules, dragees, and similar shaped, compressed pharmaceutical forms containing from about 1 ng to 500 mg of drug per unit dosage. Isotonic saline solutions can be used for parenteral administration which includes intramuscular, intrathecal, intravenous and intra-arterial routes of administration. Rectal administration can be effected through the use of suppositories formulated from conventional carriers such as cocoa butter.

Pharmaceutical compositions thus comprise one or more compounds described above and are associated with at least one pharmaceutically acceptable carrier, diluent or excipient. In preparing such compositions, the active ingredients are usually mixed with or diluted by an excipient or enclosed within such a carrier which can be in the form of a capsule or sachet. When the excipient serves as a diluent, it may be a solid, semi-solid, or liquid material which acts as a vehicle, carrier, or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, elixirs, suspensions, emulsions, solutions, syrups, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders. Examples of suitable excipients, include but are not limited to lactose, dextrose, sucrose, sorbitol, mannitol, starch, gum acacia, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose, the formulations can additionally include lubricating agents such as talc, magnesium stearate and mineral oil, wetting agents, emulsifying and suspending agents, preserving agents such as methyl- and propylhydroxybenzoates, sweetening agents or flavoring agents.

The compositions preferably are formulated in unit dosage form, meaning physically discrete units suitable as a unitary dosage, or a predetermined fraction of a unitary dose to be administered in a single or multiple dosage regimen to human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with a suitable pharmaceutical excipient. The compositions can be formulated so as to provide an immediate, sustained or delayed release of active ingredient after administration to the patient by employing procedures well known in the art.

The pharmaceutical compositions of the present invention comprise an antibiotic compound as the active ingredient, or a pharmaceutically acceptable salt thereof, and may also contain a pharmaceutically acceptable carrier, and optionally, other therapeutic ingredients, for example antivirals. The term "pharmaceutically acceptable salts" refers to

salts prepared from pharmaceutically acceptable non-toxic acids and bases, including inorganic and organic acids and bases.

The compounds of the invention that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. Acids that can be used to
5 prepare pharmaceutically acceptable acid addition salts of such basic compounds of the invention are those that form non-toxic acid addition salts, *i.e.*, salts containing pharmacologically acceptable anions, such as, but not limited to, hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, formate, acetate, propionate, succinate, camphorsulfonate, citrate, acid citrate, fumarate, gluconate,
10 isethionate, lactate, malate, mucate, gentisate, isonicotinate, saccharate, tartrate, bitartrate, para-toluenesulfonate, glycolate, glucuronate, maleate, furoate, glutamate, ascorbate, benzoate, anthranilate, salicylate, phenylacetate, mandelate, embonate (pamoate), methanesulfonate, ethanesulfonate, pantothenate, benzenesulfonate, stearate, sulfanilate, alginate, p-toluenesulfonate, and galacturonate. Particularly preferred anions are
15 hydrobromide, hydrochloride, phosphate, acid phosphate, maleate, sulfate, and acid phosphate. Most particularly preferred anions are hydrochloride and maleate.

Compounds of the invention that are acidic in nature are capable of forming salts with various pharmaceutically acceptable bases. The bases that can be used to prepare pharmaceutically acceptable base addition salts of such acidic compounds of the invention
20 are those that form non-toxic base addition salts, *i.e.*, salts containing pharmacologically acceptable cations such as, but not limited to, alkali metal or alkaline earth metal salts and the calcium, magnesium, sodium or potassium salts in particular. Suitable organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), lysine, and procaine.
25

Administration

For administration to subjects, antibiotic compounds of the invention may be formulated in pharmaceutically acceptable compositions. The compositions can be used alone or in combination with one another, or in combination with other therapeutic or
30 diagnostic agents. These compositions can be utilized in vivo, ordinarily in a mammal, preferably in a human, or in vitro. In employing them in vivo, the compositions can be administered to the mammal in a variety of ways, including parenterally, intravenously, subcutaneously, intramuscularly, colonially, rectally, vaginally, nasally, orally, transdermally, topically, ocularly, or intraperitoneally.

35

As will be readily apparent to one skilled in the art, the magnitude of a therapeutic dose of an antibiotic compound in the acute or chronic management of an infectious disease will vary with the severity of the infection or disease to be treated, the particular composition employed, and the route of administration. The dose and dose frequency will also vary according to the species of animal, age, body weight, condition and response of the individual subject. Such dosing schemes can be readily selected by those skilled in the art with due consideration of such factors.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques.

In addition to the common dosage forms set out above, an active ingredient can also be administered by controlled release means or delivery devices that are well known to those of ordinary skill in the art, such as those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; 4,008,719; 5,674,533; 5,059,595; 5,591,767; 5,120,548; 5,073,543; 5,639,476; 5,354,556; and 5,733,566, the disclosures of which are incorporated herein by reference. These dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, or microspheres or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the pharmaceutical compositions of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include: 1) extended activity of the drug; 2) reduced dosage frequency; and 3) increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and thus can affect the occurrence of side effects.

Most controlled-release formulations are designed to initially release an amount of drug that promptly produces the desired therapeutic effect, and gradually and continually

release of other amounts of drug to maintain this level of therapeutic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various inducers, including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

Pharmaceutical compositions of the invention suitable for oral administration can be presented as discrete dosage forms, such as capsules, cachets, or tablets, or aerosol sprays each containing a predetermined amount of an active ingredient as a powder or in granules, a solution, or a suspension in an aqueous or non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil liquid emulsion. Such dosage forms can be prepared by any of the methods of pharmacy, but all methods include the step of bringing the active ingredient into association with the carrier, which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation.

For example, a tablet can be prepared by compression or molding, optionally with one or more accessory ingredients. Compressed tablets can be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as powder or granules, optionally mixed with an excipient such as, but not limited to, a binder, a lubricant, an inert diluent, and/or a surface active or dispersing agent. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (*e.g.*, ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (*e.g.*, Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, for example, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, and AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PATENT, U.S.A.). An exemplary suitable binder is a mixture of microcrystalline cellulose

35

and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103™ and Starch 1500 LM.

Examples of suitable fillers for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (*e.g.*,
5 granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder/filler in pharmaceutical compositions of the present invention is typically present in about 50 to about 99 weight percent of the pharmaceutical composition.

Disintegrants are used in the compositions of the invention to provide tablets that
10 disintegrate when exposed to an aqueous environment. Too much of a disintegrant will produce tablets which may disintegrate in the bottle. Too little may be insufficient for disintegration to occur and may thus alter the rate and extent of release of the active ingredient(s) from the dosage form. Thus, a sufficient amount of disintegrant that is neither too little nor too much to detrimentally alter the release of the active ingredient(s) should be
15 used to form the dosage forms of the compounds disclosed herein. The amount of disintegrant used varies based upon the type of formulation and mode of administration, and is readily discernible to those of ordinary skill in the art. Typically, about 0.5 to about 15 weight percent of disintegrant, preferably about 1 to about 5 weight percent of disintegrant, can be used in the pharmaceutical composition.

20 Disintegrants that can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrilin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums or mixtures thereof.

25 Lubricants which can be used to form pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (*e.g.*, peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate,
30 ethyl oleate, ethyl laureate, agar, or mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL 200, manufactured by W.R. Grace Co. of Baltimore, MD), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, Texas), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, Mass), or mixtures thereof. A lubricant can optionally be added, typically in an amount of less than
35 about 1 weight percent of the pharmaceutical composition.

The methods of the invention also encompass combination therapy in which an antimicrobial compound of the invention or identified by means of the invention is administered as an admixture or sequentially with at least one other antimicrobial. The second antibacterial compound may be naturally occurring or synthetic. Suitable naturally occurring antibacterial compounds include, but are not limited to, aminoglycosides (including but not limited to dihydrostreptomycin, gentamycin, kanamycin, neomycin, paromycin and streptomycin); amphenicols (including but not limited to chloramphenicol); ansamycins (including but not limited to rifamycin); -lactams such as carbapems (including but not limited to imipenem), cephalosporins (including but not limited to cefazedone and cefroxadine), cephamycins (including but not limited to cefbuperazone); monobactams (including but not limited to aztreonam), oxacephems (including but not limited to flomoxef) or penicillins (including but not limited to ampicillin, carbencillin, methicillin, penicillin N, penicillin O and penicillin V); lincosamides (including but not limited to clindamycin and lincomycin); macrolides (including but not limited to carbomycin and erythromycin); polypeptides (including but not limited to gramicidin S); glycopolypeptides (including but not limited to vancomycin and teichoplanin); tetracyclines (including but not limited to apicycline, methacycline and tetracycline); and others such as cycloserine, mupirocin and tuberin. Suitable synthetic antibacterial compounds include 2,4-diaminopyrimidines (including but not limited to trimethoprim); nitrofurans (including but not limited to nifuradene); quinolones and quinolone analogs (including but not limited to enoxacin, lomefloxacin, nalidixic acid and ofloxacin); streptogramins; sulfonamides (including but not limited to sulfamoxole and sulfanilamide); sulfones (including but not limited to diathymosulfone); oxazolidinones (including but not limited to linezolid); and others such as glycylcyclines, clofoctol, hexedine, methenamine, and nitroxoline.

The "adjunct administration" of a compound identified by the method of the invention and a second antibacterial compound means that the two are administered either as a mixture or sequentially. When administered sequentially, the compound may be administered before or after the second antibacterial compound, so long as the initially administered compound is still providing antibacterial activity. Any of the above described modes of administration can be used in combination to deliver the compound and the second antibacterial compound. When a compound identified by the method of the invention and a second antibacterial compound are administered adjunctively as a mixture, they are preferably given in the form of a pharmaceutical composition comprising both agents. Thus, in a further embodiment of the invention, it is provided a pharmaceutical

35

composition comprising a compound of the invention and a second antibacterial compound together with a pharmaceutically acceptable carrier.

Target Infectious Agents

5 The antibiotic compounds identified by the methods of the invention can be used to treat infectious diseases in animals, including humans, companion animals (*e.g.*, dogs and cats), livestock animals (*e.g.*, sheep, cattle, goats, pigs, and horses), laboratory animals (*e.g.*, mice, rats, and rabbits), and captive or wild animals.

10 The antibiotics of the invention selectively target the invading microorganism and thus cause minimal adverse reactions common to many antibiotics such as gastrointestinal disturbances, allergies and hypersensitivities, blood disorders and central nervous system toxicities. The compounds of the invention are expected to be particularly effective against pathogenic organisms found in the blood, muscle tissue (including heart), urinary tract, lower and upper respiratory tract, intestinal, skin and wound, genital tract, and mucosal

15 tissue and include peripheral and central nervous system tissue but are not limited to, gram positive cocci, such as Staphylococci (*e.g.*, *S. aureus*), Streptococci (*e.g.*, *S. pneumoniae*, *S. pyrogens*, *S. faecalis*, *S. viridans*); gram positive bacilli, such as *Bacillus* (*e.g.*, *B. anthracis*), *Corynebacterium* (*e.g.*, *C. diphtheriae*), *Listeria* (*e.g.*, *L. monocytogenes*); gram negative cocci, such as *Neisseria* (*e.g.*, *N. gonorrhoeae*, *N. Meningitidis*); gram negative bacilli, such

20 as *Haemophilus* (*e.g.*, *H. influenzae*), *Pasteurella* (*e.g.*, *P. multocida*), *Proteus* (*e.g.*, *P. mirabilis*), *Salmonella* (*e.g.*, *S. typhimurium*), *Shigella* species, *Escherichia* (*e.g.*, *E. coli*), *Klebsiella* (*e.g.*, *K. pneumoniae*), *Serratia* (*e.g.*, *S. marcescens*), *Yersinia* (*e.g.*, *Y. pestis*), *Providencia* species, *Enterobacter* species, *Bacteroides* (*e.g.*, *fragilis*), *Acinetobacter* species, *Campylobacter* (*e.g.*, *C. jejuni*), *Pseudomonas* (*e.g.*, *P. aeruginosa*), *Bordetella* (*e.g.*

25 , *B. pertussis*), *Brucella* species, *Fracisella* (*e.g.*, *F. tularensis*), *Clostridia* (*e.g.*, *C. perfringens*), *Helicobacter* (*e.g.*, *H. pylori*), *Vibrio* (*e.g.*, *V. cholerae*), *Mycoplasma* (*e.g.*, *M. pneumoniae*), *Legionella* (*e.g.*, *L. pneumophila*), *Spirochetes* (*e.g.*, *Treponema*, *Leptospira* and *Borrelia*), *Mycobacteria* (*e.g.*, *M. tuberculosis*), *Nocardia* (*e.g.*, *N. asteroides*), *Chlamydia* (*e.g.*, *C. trachomatis*), and *Rickettsia* species.

30 The following examples serve to further typify the nature of the invention but should not be construed as a limitation in the scope thereof, which scope is defined solely by the appended claims.

5. EXAMPLES

The compounds of the invention can be prepared using commercially available starting materials and commercially available reagents using a variety of starting organic transformations. Nevertheless, specific synthetic routes and examples of syntheses are disclosed herein for illustrative purposes.

General

¹H and ¹³C NMR spectra were recorded on a Varian 300 MHz FT-NMR.

Preparative HPLC purification was performed on a Gilson HPLC System with 215 autosampler/fraction collector using a Hewlett-Packard 50 x 21.2 mm XDB C18 reverse-phase column running acetonitrile/water gradients at 25 ml/min. Analytical HPLC/MS (electrospray) was performed on a Hewlett-Packard 1100 HPLC with MSD using a MetaChem MetaSil AQ C18 column running acetonitrile/water gradients at 2 mL/min. All anhydrous solvents were purchased from Aldrich chemical company in 15 SureSeal containers. Most reagents were purchase from Aldrich Chemical Company. Protected amino acids were obtained from NovaBiochem.

Abbreviations: Boc, tert-butoxycarbonyl; DCE, 1,2-dichloroethane; DCM, dichloromethane; TEA, triethylamine; HOBt, 1-hydroxybenzotriazole; DMAP, dimethylaminopyridine; EDC, 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide 20 hydrochloride; NHS, N-Hydroxysuccinimide; DMF, dimethylformamide; EtOAc, ethyl acetate; MeOH, methanol; OSu N-hydroxysuccinamide ester; THF, tetrahydrofuran; TFA, trifluoroacetic acid.


Array synthesis was conducted in 15 x 75 mm glass round bottom screw-cap vials contained in a custom 4 x 6 array aluminum synthesis block and sealed with a Teflon-lined 25 rubber membrane. Reagents were added and aqueous extractions performed with single or multichannel pipettors. Filtrations were performed using Whatman/Polyfiltronics 24 well, 10 mL filtration blocks on a Whatman/Polyfiltronics UniVac filtration manifold modified to collect eluent in the custom synthesis block. Evaporation of volatile materials from the array was performed with a Labconco Vortex-Evaporator.

30

5.1.1 Synthesis of Pyrrolidine Carbamate Series(I)

Compounds represented by Formula VI were synthesized in arrays as described in section 5.1.1. See Scheme I for a general synthetic scheme and Table 1 for representative R groups. The general synthetic scheme involved initial treatment of tert-butyl-(2-pentyl) 35 succinate-mono-N-hydroxysuccinimide ester with 3-pyrrolidinol to afford the

3-hydroxypyrrolidine amide. The tert-butyl group was then removed with subsequent conversion of the carboxylic acid to the methyl ester. The esters were treated with DMAP and various isocyanates were utilized to convert the alcohol to the pyrrolidine carbamates. Finally, the ester moiety was treated with a methanolic solution of hydroxylamine hydrochloride to yield the hydroxamic acid.



Formula VI

Step 1:

To a solution of (R) tert-butyl (2-pentyl)succinate mono N-hydroxysuccinimide ester (1 g, 2.9 mmol) in DCM was added 3-pyrrolidinol (0.51 g, 5.8 mmol). The resulting mixture was stirred at room temperature overnight and evaporated to dryness. The residue was dissolved in EtOAc and washed with 1 N HCl followed by brine. The organic layer was dried over anhydrous sodium sulfate, filtered and solvent removed under reduced pressure. The crude product was purified by column chromatography using EtOAc as eluting solvent. Yield 0.62 g, 68%. ¹H NMR (300 MHz, CDCl₃) δ: 0.95 (t, 3H), 1.30 (m, 6H), 1.49 (s, 9H), 1.51 (m, 1H), 1.68 (m, 1H), 2.11 (m, 1H), 2.13 (m, 1H), 2.34 (m, 1H), 2.77-3.01 (m, 3H), 3.50-3.80 (m, 3H), 3.95 (m, 1H), 4.56 (m, 1H). HPLC/MS calculated weight 313.44; found 314.4, 335.4 (M+H, M+Na).

Step 2:


The t-butyl ester (620 mg, 1.98 mmol) was subjected to 10 ml of 20% TFA/DCM solution for 4 h. The solvent was evaporated and dried under high vacuum to yield crude acid. The acid was dissolved in 3 ml of methanol and 6 ml of benzene, and trimethylsilyldiazomethane in hexanes (2.0 M, 6.0 ml, 12 mmol) was added slowly. Volatile material was removed under vacuum and the residue was purified by flash chromatography using 2% MeOH/EtOAc. Yield 480 mg, 90%. ¹H NMR (300 MHz, CDCl₃) δ: 0.96 (t, 3H), 1.35 (m, 6H), 1.52 (m, 1H), 1.70 (m, 1H), 2.01-2.19 (m, 3H), 2.52 (m, 1H), 2.89-3.04 (m, 2H), 3.55-3.72 (m, 2H), 3.72 (s, 3H), 3.78 (m, 1H), 3.96 (m, 1H), 4.58 (m, 1H). HPLC/MS calculated weight 271.36; found 272.4, 294.4 (M+H, M+Na).

Step 3:

To each tube in a 4 x 6 array containing the alcohol ester from the previous reaction (32.5 mg, 0.12 mmol), TEA (33 ml, 0.24 mmol), and DMAP (5 mg) in 2 ml of THF was added an isocyanate (0.36 mmol). The mixtures were agitated and heated at 80°C for 4 h then at 60°C for 12 h, evaporating to dryness. The residue was brought up in 4 ml DCM then washed with 0.5 N aqueous HCl (2 ml) and saturated aqueous NaCl (2 ml). The organic layers were filtered through approximately 0.5 g of anhydrous Na₂SO₄ each and evaporated to dryness. The resulting pyrrolidine carbamates (oils) were each placed in a vacuum chamber under high vacuum overnight, weighed, and analyzed by HPLC/MS.

Step 4:

A methanolic solution of NH₂OH.HCl (2 M, 30 ml) was cooled to 0°C. A methanolic solution of KOH (3 M, 30 ml) was added, the mixture was stirred for 30 min and filtered. An aliquot (2 ml, 2.0 mmol NH₂OH) of this solution was added to each of the reaction vials containing the methyl ester product of the previous reaction. The array was agitated for 30 min. Each solution was acidified to pH = 5-6 with Dowex H⁺ resin. The solutions were then filtered and concentrated to dryness. The crude materials were dissolved in 1 ml of DMSO and purified by preparative HPLC. The product-containing fractions were collected, evaporated in a Savant evaporating centrifuge, and re-analyzed by HPLC/MS.



Scheme I

The following compounds were synthesized by the procedure noted in section 5.1.1.

The following isocyanates were used: 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 2-fluoro-phenyl, 3-fluoro-phenyl, 4-fluoro-phenyl, 2-bromo-phenyl, 3-bromo-phenyl,

phenyl, 4-methoxy-phenyl, 4-trifluoro-methoxy-phenyl, 4-(N,N-Dimethyl-amino)-phenyl,
4-bromo-phenyl, 4-methyl-phenyl, 4-n-butyl-phenyl, 4-trifluoro-methyl-phenyl,
2-methoxy-phenyl, 2-ethyl-phenyl, 2-i-propyl-phenyl, 3-methyl-phenyl, 3-nitro-phenyl,
3-methyl-mercapto-phenyl, 4-chloro-3-nitro-phenyl, 4-chloro-3-trifluoromethyl-phenyl,
5 2,4-dichloro-phenyl, 2,4-dimethoxy-phenyl, 4-chloro-2-tri-fluoromethyl-phenyl,
2,5-difluoro-phenyl, 2-methoxy-5-chloro-phenyl, 2-methoxy-5-methyl-phenyl,
3,5-dimethyl-phenyl, 3,5-bis-(tri-fluoromethyl)-phenyl, 3,5-dichloro-phenyl,
4-(2,6-dichloro-pyridyl), n-propyl, benzyl, 2-(ethyl-2-thiophene), 4-phenoxy-phenyl,
4-trifluoro-methanemercapto-phenyl, 2-phenyl-phenyl, 2-phenoxy-phenyl.

10 All compounds were analyzed by HPLC/MS.

Synthesis of compound 1. Calculated weight 425.91, Found 394.2, 427.2, 449.2
(M-NHOH, M+H, M+Na).

15 **Synthesis of compound 2.** Calculated weight 425.91, Found 394.2, 427.2, 449.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 3. Calculated weight 425.91, Found 394.2, 427.2, 449.2
(M-NHOH, M+H, M+Na).

20

Synthesis of compound 4. Calculated weight 409.46, Found 377.4, 410.2, 432.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 5. Calculated weight 409.46, Found 377.4, 410.2, 432.2
25 (M-NHOH, M+H, M+Na).

Synthesis of compound 6. Calculated weight 409.46, Found 377.4, 410.2, 432.2
(M-NHOH, M+H, M+Na).

30 **Synthesis of compound 7.** Calculated weight 470.36, Found 438.0, 471.0, 493.0
(M-NHOH, M+H, M+Na).

Synthesis of compound 8. Calculated weight 470.36, Found 471.0, 493.0 (M+H,
M+Na).

35

Synthesis of compound 9. Calculated weight 391.47, Found 359.2, 392.3, 414.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 10. Calculated weight 421.49, Found 389.2, 422.2, 444.2
5 (M-NHOH, M+H, M+Na).

Synthesis of compound 11. Calculated weight 475.47, Found 443.2, 476.2, 498.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 12. Calculated weight 434.54, Found 402.4, 435.3, 457.2
10 (M-NHOH, M+H, M+Na).

Synthesis of compound 13. Calculated weight 470.36, Found 438.0, 471.0, 493.0
(M-NHOH, M+H, M+Na).
15

Synthesis of compound 14. Calculated weight 405.49, Found 373.3, 406.2, 428.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 15. Calculated weight 447.58, Found 415.3, 448.2, 470.2
20 (M-NHOH, M+H, M+Na).

Synthesis of compound 16. Calculated weight 459.47, Found 427.1, 482.0
(M-NHOH, M+Na).

Synthesis of compound 17. Calculated weight 421.49, Found 389.2, 422.2, 444.2
25 (M-NHOH, M+H, M+Na).

Synthesis of compound 18. Calculated weight 419.52, Found 387.2, 420.2, 442.3
(M-NHOH, M+H, M+Na).
30

Synthesis of compound 19. Calculated weight 433.55, Found 401.2, 434.2, 456.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 20. Calculated weight 405.49, Found 373.3, 406.2, 428.3
35 (M-NHOH, M+H, M+Na).

Synthesis of compound 21. Calculated weight 436.47, Found 404.0, 437.3, 459.2
(M-NHOH, M+H, M+Na).

5 **Synthesis of compound 22.** Calculated weight 437.56, Found 405.2, 438.3, 460.1
(M-NHOH, M+H, M+Na).

Synthesis of compound 23. Calculated weight 470.91, Found 439.2, 472.2, 494.2
(M-NHOH, M+H, M+Na).

10 **Synthesis of compound 24.** Calculated weight 493.91, Found 462.0, 495.0, 517.0
(M-NHOH, M+H, M+Na).

Synthesis of compound 25. Calculated weight 460.36, Found 428.0, 461.0, 483.0
(M-NHOH, M+H, M+Na).

15 **Synthesis of compound 26.** Calculated weight 451.52, Found 419.2, 452.2, 474.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 27. Calculated weight 493.91, Found 462.0, 495.0, 517.0
20 (M-NHOH, M+H, M+Na).

Synthesis of compound 28. Calculated weight 427.45, Found 395.2, 428.3, 450.1
(M-NHOH, M+H, M+Na).

25 **Synthesis of compound 29.** Calculated weight 455.94, Found 423.1, 457.1, 479.1
(M-NHOH, M+H, M+Na).

Synthesis of compound 30. Calculated weight 435.52, Found 403.3, 436.2, 458.3
(M-NHOH, M+H, M+Na).

30 **Synthesis of compound 31.** Calculated weight 419.52, Found 387.2, 420.2, 442.3
(M-NHOH, M+H, M+Na).

Synthesis of compound 32. Calculated weight 527.46, Found 495.0, 528.3, 550.0
35 (M-NHOH, M+H, M+Na).

Synthesis of compound 33. Calculated weight 460.36, Found 428.0, 461.0, 483.0
(M-NHOH, M+H, M+Na).

Synthesis of compound 34. Calculated weight 461.34, Found 429.0, 462.0, 483.0
5 (M-NHOH, M+H, M+Na).

Synthesis of compound 35. Calculated weight 357.45, Found 325.4, 358.4, 380.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 36. Calculated weight 405.49, Found 373.4, 406.4, 428.3
10 (M-NHOH, M+H, M+Na).

Synthesis of compound 37. Calculated weight 425.55, Found 393.2, 426.2, 448.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 38. Calculated weight 483.56, Found 451.2, 484.2, 506.2
15 (M-NHOH, M+H, M+Na).

Synthesis of compound 39. Calculated weight 491.53, Found 459.0, 492.0, 514.1
20 (M-NHOH, M+H, M+Na).


Synthesis of compound 40. Calculated weight 467.57, Found 435.2, 468.2, 490.2
(M-NHOH, M+H, M+Na).

Synthesis of compound 41. Calculated weight 483.56, Found 451.2, 484.2, 506.1
25 (M-NHOH, M+H, M+Na).

5.1.2 Synthesis of Pyrrolidine Carbamate Series(II)

Compounds represented by Formula VII were synthesized as described in section
30 5.1.2. See Scheme I for a general synthetic scheme and Table 2 for representative R groups.
Synthesis of Series II of the pyrrolidine carbamates was accomplished in an analogous way
to Series I, with the exception that in Step 3 secondary amines were used instead of
isocyanates.

35



Formula VII

Step 1:

As in step 1 of the pyrrolidine carbamate I series, section 5.1.1.

Step 2:

As in step 2 of the pyrrolidine carbamate I series, section 5.1.1.

Step 3:

To a solution of triphosgene (78 mg, 0.29 mmol) in anhydrous DCM (2 ml) at 0°C was added the alcohol from Step 2 (78 mg, 0.29 mmol) followed by the addition of TEA (120 ml, 0.87 mmol). The ice bath was removed and resulting mixture was allowed to stir for 4 h. Volatile material was removed in vacuo and residue was taken up in 2 ml of anhydrous DCM. To this solution was added secondary amine (1.44 mmol) and the mixture was allowed to shake for 12 h. The solution was diluted with 2 ml of DCM and washed with 5% aqueous citric acid and brine. The organic layer was dried over sodium sulfate and concentrated to dryness. The crude residue was used in the next step without further purification.

Step 4:

As in step 4 of the pyrrolidine carbamate I series, section 5.1.1.

The following secondary amines were used: N-pyrrolidine, N-pyrrolidine-2-methanol, N-piperidine, N-homo-piperidine, N-morpholine, N-3S-pyrrolidinol

Synthesis of compound 42. Calculated weight 369.46, Found 337.4, 370.4, 392.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 43. Calculated weight 399.49, Found 367.3, 400.4, 422.4 (M-NHOH, M+H, M+Na).

Synthesis of compound 44. Calculated weight 399.49, Found 367.3, 400.3, 422.4 (M-NHOH, M+H, M+Na).

Synthesis of compound 45. Calculated weight 383.49, Found 351.2, 384.2, 406.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 46. Calculated weight 397.52, Found 365.2, 398.3, 420.2 (M-NHOH, M+H, M+Na).


Synthesis of compound 47. Calculated weight 385.46, Found 353.2, 386.2, 408.3 (M-NHOH, M+H, M+Na).

Synthesis of compound 48. Calculated weight 385.46, Found 353.2, 386.2, 408.3 (M-NHOH, M+H, M+Na).

5.2.1 Synthesis of Piperidine Carbamate Series

Compounds represented by Formula X were synthesized as described in section

5.2.1. See Scheme II for general synthetic scheme and Table 3 for representative R groups. The piperidine carbamate series was synthesized analogously to the pyrrolidine series (See 5.1.1 above) replacing 3-pyrrolidinol with 3-hydroxypiperidine hydrochloride salt and proportionate amounts of the other reagents.



Formula X

Step 1:

To a solution of (R) tert-butyl (2-pentyl)succinate mono N-hydroxysuccinimide ester (2 g, 5.86 mmol) in DCM was added 3-hydroxypiperidine hydrochloride (1.6 g, 11.7 mmol) and TEA (2 ml, 14.65 mmol). The resulting mixture was stirred at room temperature overnight and evaporated to dryness. The residue was dissolved with EtOAc

and washed with 1 N HCl followed by brine. The organic layer was dried over anhydrous sodium sulfate, filtered and solvent removed under reduced pressure. The crude product was purified by column chromatography using 75% EtOAc in hexane. Yield 1.75 g, 92%.

¹H NMR (300 MHz, CDCl₃) δ: 0.95 (t, 3H), 1.34 (m, 6H), 1.50 (s, 9H), 1.57 (m, 4H), 1.86-1.97 (m, 3H), 2.45 (m, 1H), 2.91-2.97 (m, 1H), 3.20-3.23 (m, 1H), 3.49-3.54 (m, 1H), 4.02-4.24 (m, 3H), 4.47 (m, 1H). HPLC/MS calculated weight 327.47; found 328.4, 350.4 (M+H, M+Na).

Step 2:


As in Step 2 of pyrrolidine carbamate series, section 5.1.1. Yield 1.05 g, 70%. ¹H NMR (300 MHz, CDCl₃) δ: 0.95 (t, 3H), 1.34 (m, 6H), 1.51-1.68 (m, 4H), 1.86-1.97 (m, 3H), 2.46 (m, 1H), 2.91-3.03 (m, 1H), 3.20-3.26 (m, 1H), 3.49-3.61 (m, 1H), 3.74 (s, 3H), 3.89-4.03 (m, 3H), 4.37 (m, 1H). HPLC/MS calculated weight 285.39; found 286.4, 308.4 (M+H, M+Na).

Step 3:

As in step 3 of pyrrolidine carbamate series I, section 5.1.1.

Step 4:

As in step 4 of pyrrolidine carbamate series I, section 5.1.1.



Scheme II

The following isocyanates were used: phenyl, 3,5-di-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 4-methoxy-phenyl, 4-(N,N-di-methylamino)-phenyl, 4-trifluoro-methoxy-phenyl, 4-fluoro-phenyl

Synthesis of compound 49. Calculated weight 405.49, Found 373.3, 406.3, 428.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 50. Calculated weight 474.38, Found 442.2, 475.2, 497.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 51. Calculated weight 439.94, Found 408.1, 441.2, 463.1 (M-NHOH, M+H, M+Na).

Synthesis of compound 52. Calculated weight 439.94, Found 408.1, 441.1, 463.1 (M-NHOH, M+H, M+Na).

Synthesis of compound 53. Calculated weight 435.52, Found 403.3, 436.2, 458.3 (M-NHOH, M+H, M+Na).

Synthesis of compound 54. Calculated weight 448.56, Found 416.2, 449.2, 471.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 55. Calculated weight 489.49, Found 457.2, 490.2, 512.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 56. Calculated weight 423.48, Found 391.2, 424.2, 446.2 (M-NHOH, M+H, M+Na)


5.2.2 Synthesis of Tic and Pyrrolidine Series

Compounds represented in Table 5 were synthesized as described in section 5.2.2. See Scheme IIIA for general synthetic scheme and Table 5 for representative R groups.

Step 1:

To a solution of (S)-4-Benzyl-2-oxazolidione (6.0 g, 0.034 mol) in dry THF (10 ml/g) at -78°C under argon was added 13.6 ml of n-butyllithium (2.5 M) over 20 min, and the resulting mixture was stirred for 1 h at -78°C. Heptanoyl chloride (5.8 ml, 0.037 mol) was slowly added and allowed to stir at -78°C for 1h and warm to room temperature over an additional 1 h. The reaction was quenched with saturated NH₄Cl (50% total reaction volume), the THF was removed, and the aqueous phase was washed with DCM (2x). The combined organic layers were washed with 1N NaOH and brine, dried over anhydrous Na₂SO₄. The crude product was obtained after filtration and removal of solvent. The isolated product was utilized in the next step without further purification. Yield: 9.13 g,

93%. Analyzed by ^1H NMR (300MHz, CDCl_3): δ 0.82 (t, 3H), 1.27 (m, 6H), 1.59 (m, 2H), 2.69 (dd, 1H), 3.22 (m, 1H), 4.10 (m, 2H), 4.59 (m, 1H), 7.26 (m, 5H). HPLC/MS. Calculated weight 289, Found 290.3, 312.2 (M+H, M+Na).



Scheme IIIA

Step 2:

Lithium hexamethyldisilazide (118 ml, 1M in THF) was cooled to -78°C under argon and a solution of an N-heptanoyloxazolidinone (31g, 0.107 mol) in dry THF (5 ml/g) was added while maintaining the internal reaction temperature below -65°C . After addition, the resulting solution was allowed to stir at -78°C for 1 h. A solution of methyl bromoacetate (24.63g, 0.161 mol) in dry THF (1 ml/g) was added while the internal reaction temperature was maintained below -70°C . The resulting mixture was stirred at -78°C for 3 h and then allowed to warm to -40°C over an additional 1 h. The reaction mixture was then warm to 0°C and quenched with saturated NH_4Cl (50% total reaction volume). The THF was removed under vacuum, and the aqueous phase was washed with EtOAc (2x). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered, and concentrated. The residue was purified by flash chromatography (15% EtOAc/hexane) to give desired product.

Yield: 30.07g of pale yellow oil, 78%. Analyzed by : ^1H NMR (300MHz, CDCl_3): δ 0.96 (t,


3H), 1.37 (m, 6H), 1.56 (m, 1H), 1.75 (m, 1H), 2.64 (dd, 1H), 2.83 (dd, 1H), 3.00 (dd, 1H), 3.42 (dd, 1H), 3.75 (s, 3H), 4.24 (m, 3H), 4.75 (m, 1H), 7.36 (m, 5H). HPLC/MS. Calculated weight 361, Found 362.3, 384.2 (M+H, M+Na).

Step 3:

- 5 An alkylated N-heptanoyloxazolidinones (3.6 g, 0.01 mol) was dissolved in 36 ml of THF/H₂O (4/1, 10 ml/g) and cooled to 0°C. Hydrogen peroxide (30% in water, 6.1 ml, 0.62 ml/mmol) was added followed by a solution of LiOH in water (0.56 g, 0.013 mol, 1.18 M). The reaction mixture was stirred at 0°C for 3 h, after which a solution of Na₂SO₃ in water (1.67 g, 0.013 mol, 2.73 M) was added while maintaining the temperature below
- 10 10°C. The THF was removed and the remaining aqueous phase was washed with DCM (2x). The combined organic layers were washed with saturated aqueous NaHCO₃ (2x). The aqueous phase was then acidified to pH = 3 with 1N HCl and washed with EtOAc (3x). The combined organic extracts were dried over anhydrous Na₂SO₄, filtered, and concentrated to give desired product. Yield 1.5 g, 75%. Analyzed by ¹H NMR (300MHz,
- 15 CDCl₃): δ 0.97 (t, 3H), 1.38 (m, 6H), 1.65 (m, 1H), 1.77 (m, 1H), 2.54 (dd, 1H), 2.80(dd, 1H), 2.96 (m, 1H), 3.78 (s, 3H). HPLC/MS. Calculated weight 202, Found 225.1 (M+Na).

5.2.3 Synthesis of Thioproline Series

- Compounds represented by Formula VIII were synthesized as described in section
- 20 5.2.2. See Scheme III for a general synthetic scheme and Table 4 for representative R groups. The thioproline series was synthesized with a solution of EDC and HOBT Boc-thioproline and an amine to afford the amino acid amide. To a solution of 2-methylaceto-hexanoic acid with HOBT and EDC was added the amino acid amide. The resulting residue was subsequently treated with hydroxylamine to afford the desired
- 25 product.



Formula VIII

Synthesis of compound 57.**Step 1 and 2 (Scheme III):**


1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (0.290 g, 1.52 mmol) and HOBt (0.256 g, 1.89 mmol) were added to a solution of Boc-thiopropine (0.294 g, 1.26 mmol) in DMF (3.75 ml) and agitated for 1 h. Isobutylamine (0.159 ml, 1.6 mmol) was added and the solution was agitated overnight. Saturated ammonium chloride solution was added (5 mL) and the mixture was extracted with EtOAc. The organic layer was washed with saturated NaHCO_3 , 0.5 N HCl, followed by brine and dried over Na_2SO_4 . The solvent was removed in vacuo to give crude Boc aminoamide. The Boc group was removed by treatment with 2 ml of 4.0 M HCl/dioxane for 1 h followed by removal of the solvent in vacuo to give the HCl salt of the amino amide.

Step 3 (Scheme III):

The amino amide HCl salt from Step 1 was added along with triethylamine (0.139 ml, 1.00 mmol) to a solution of (R)-methyl-(2-pentyl)succinate monoacid (0.050 g, 0.25 mmol), EDC (0.057 g, 0.30 mmol) and HOBt (0.50 g, 0.37 mmol) which had been agitated for 30 min. The resulting mixture was allowed to shake overnight. A saturated solution of ammonium chloride (2.5 ml) was added and the mixture was extracted with EtOAc. The organic layer was washed with saturated NaHCO_3 , 0.5 N HCl, followed by brine and dried over Na_2SO_4 . The solvent was removed in vacuo.

Step 4 (Scheme III):

The residue from Step 3 was dissolved in 2 ml of a solution prepared by mixing equal parts of cold 3 M KOH in methanol and cold 2 M NH_2OH and filtering. This was allowed to shake for 15 min and Dowex H⁺ resin was added to reach a pH ~ 7. The solution was filtered, concentrated and purified by reverse phase HPLC to give 5.5 mg of Compound 57. Analyzed by HPLC/MS. Calculated weight 373.52, Found 341.1, 374.2, 396.1 (M-NHOH, M+H, M+Na).



Scheme III

Synthesis of compound 58. Synthesis of compound 58 was completed as described above for compound 57 using cyclopentyl amine in place of isobutylamine. Calculated weight 385.53, Found 353.2, 386.2, 408.1 (M-NHOH, M+H, M+Na).

5 **Synthesis of compound 59.** Synthesis of compound 59 was completed as described above for compound 57 using cyclohexylamine in place of isobutylamine. Calculated weight 399.55, Found 367.3, 400.2 (M-NHOH, M+H).

10 **Synthesis of compound 60.** Synthesis of compound 60 was completed as described above for compound 57 using tetrahydrofurfuryl-amine in place of isobutylamine. Calculated weight 401.53, Found 369.2, 402.2, 424.2 (M-NHOH, M+H, M+Na).

15 **Synthesis of compound 61.** Synthesis of compound 61 was completed as described above for compound 57 using furfurylamine in place of isobutylamine. Calculated weight 39.50, Found 365.1, 398.1, 420.0 (M-NHOH, M+H, M+Na).

20 **Synthesis of compound 62.** Synthesis of compound 62 was completed as described above for compound 57 using 2-thiophenmethylamine in place of isobutylamine. Calculated weight 413.56, Found 381.0, 414.1, 436.0 (M-NHOH, M+H, M+Na).

25 **Synthesis of compound 63.** Synthesis of compound 63 was completed as described above for compound 57 using benzylamine in place of isobutylamine. Calculated weight 407.53, Found 375.2, 408.3, 430.1 (M-NHOH, M+H, M+Na).

30 **Synthesis of compound 64.** Synthesis of compound 64 was completed as described above for compound 57 using 2-aminothiazole in place of isobutylamine. Calculated weight 400.52, Found 368.0, 401.0, 423.0 (M-NHOH, M+H, M+Na).

35 **Synthesis of compound 65.** Synthesis of compound 65 was completed as described above for compound 57 using dimethylamine in place of isobutylamine. Calculated weight 345.46, Found 313.2, 346.3, 368.1 (M-NHOH, M+H, M+Na).

Synthesis of compound 66. Synthesis of compound 66 was completed as described above for compound 57 using pyrrolidine in place of isobutylamine. Calculated weight 371.50, Found 339.2, 372.2, 394.1 (M-NHOH, M+H, M+Na).

Synthesis of compound 67. Synthesis of compound 67 was completed as described above for compound 57 using piperidine in place of isobutylamine. Calculated weight 385.53, Found 353.2, 386.2, 408.1 (M-NHOH, M+H, M+Na).

5 **Synthesis of compound 68.** Synthesis of compound 68 was completed as described above for compound 57 using hexamethylineimine in place of isobutylamine. Calculated weight 399.55, Found 367.3, 400.2, 422.1 (M-NHOH, M+H, M+Na).

10 **Synthesis of compound 69.** Synthesis of compound 69 was completed as described above for compound 57 using morpholine in place of isobutylamine. Calculated weight 387.50, Found 355.2, 388.2, 410.1 (M-NHOH, M+H, M+Na).


15 **Synthesis of compound 70.** Synthesis of compound 70 was completed as described above for compound 57 using 2,6-dimethylmorpholine in place of isobutylamine. Calculated weight 415.55, Found 383.2, 416.2, 438.1 (M-NHOH, M+H, M+Na).

20 **Synthesis of compound 71.** Synthesis of compound 71 was completed as described above for compound 57 using thiomorpholine in place of isobutylamine. Calculated weight 403.57, Found 371.0, 404.2, 426.0 (M-NHOH, M+H, M+Na).

5.2.3 Synthesis of Tic and Proline Series


Compounds of the formulas listed below were synthesized in a manner similar to the thioproline series. See section 5.2.2 and Table 5 for representative R groups.

25 **Synthesis of compound 72.** Synthesis of compound 72 was completed as described above for compound 57 using Boc-D-tetrahydroisoquinoline in place of Boc-thioproline. Calculated weight 415.54, Found 345.2, 416.2 (M-pyrrolidine, M+H).




Compound 72

Synthesis of compound 73. Synthesis of compound 73 was completed as described above for compound 57 using Boc-tetrahydroisoquinoline in place of Boc-thioproline. Calculated weight 415.54, Found 345.1, 438.3 (M-pyrrolidine, M+Na).




Compound 73

Synthesis of compound 74. Synthesis of compound 74 was completed as described above for compound 57 using Boc-L-proline in place of Boc-thioproline. Calculated weight 353.47, Found 321.2, 354.2 (M-NHOH, M+H).



Compound 74


Synthesis of compound 75. Synthesis of compound 75 was completed as described above for compound 57 using Boc-D-proline in place of Boc-thioproline. Calculated weight 353.47, Found 321.2, 354.2 (M-NHOH, M+H).



Compound 75

5.2.4 Synthesis of Benzyloxypoline Series

Compounds of the benzyloxypoline series, Formula IX were synthesized in a manner similar to the thiopoline series. See section 5.2.3 and Table 6 for representative R groups.



Formula IX

Synthesis of compound 76. Synthesis of compound 76 was completed as described above for compound 64 using Boc-3-benzyloxypoline in place of Boc-thiopoline.

Calculated weight 488.61, Found 456.1; 489.2 (M-NHOH, M+H).

Synthesis of compound 77. Synthesis of compound 77 was completed as described above for compound 65 using Boc-3-benzyloxypoline in place of Boc-thiopoline.

Calculated weight 433.55, Found 401.2, 434.2 (M-NHOH, M+H).

Synthesis of compound 78. Synthesis of compound 78 was completed as described above for compound 87 using Boc-3-benzyloxypoline in place of Boc-homopoline.

Calculated weight 495.62, Found 463.3, 496.3 (M-NHOH, M+H).

Synthesis of compound 79. Synthesis of compound 79 was completed as described above for compound 58 using Boc-3-benzyloxypoline in place of Boc-thiopoline.


Calculated weight 473.62, Found 441.2, 474.2 (M-NHOH, M+H).

Synthesis of compound 80. Synthesis of compound 80 was completed as described above for compound 71 using Boc-3-benzyloxypoline in place of Boc-thiopoline.

Calculated weight 491.65, Found 469.3, 492.3 (M-NHOH, M+H).


5.2.5 Synthesis of Homopoline Series I

Compounds represented by Formula X are synthesized as described in section 5.2.2. See Scheme IV for a general synthetic scheme and Table 7 for representative R groups. The



Scheme IV

homoproline series was synthesized in an analogous fashion to the thioproline series, with Boc-thioproline replaced by Boc-pipecolic acid.



Formula X

Synthesis of compound 81. Synthesis of compound 81 was completed as described above for compound 57 using Boc-pipecolic acid in place of Boc-thioproline.
 20 Calculated weight 369.50, Found 337.2, 392.3 (M-NHOH, M+Na).

Synthesis of compound 82. Synthesis of compound 82 was completed as described above for compound 58 using Boc-pipecolic acid in place of Boc-thioproline.
 Calculated weight 381.52, Found 349.2, 404.3 (M-NHOH, M+Na).
 25

Synthesis of compound 83. Synthesis of compound 83 was completed as described above for compound 59 using Boc-pipecolic acid in place of Boc-thioproline.
 Calculated weight 395.54, Found 363.3, 418.2 (M-NHOH, M+Na).

30 **Synthesis of compound 84.** Synthesis of compound 84 was completed as described above for compound 60 using Boc-pipecolic acid in place of Boc-thioproline.
 Calculated weight 397.52, Found 365.2, 420.0 (M-NHOH, M+Na).

35

Synthesis of compound 85. Synthesis of compound 85 was completed as described above for compound 61 using Boc-pipecolic acid in place of Boc-thioprolin. Calculated weight 393.48, Found 361.2, 416.2 (M-NHOH, M+Na).

5 **Synthesis of compound 86.** Synthesis of compound 86 was completed as described above for compound 62 using Boc-pipecolic acid in place of Boc-thioprolin. Calculated weight 409.55, Found 377.2, 432.2 (M-NHOH, M+Na).

10 **Synthesis of compound 87.** Synthesis of compound 87 was completed as described above for compound 57 using Boc-pipecolic acid in place of Boc-thioprolin and 2-aminomethylpyridine in place of isobutylamine. Calculated weight 404.51, Found 405.2, 427.2 (M+H, M+Na).

15 **Synthesis of compound 88.** Synthesis of compound 88 was completed as described above for compound 63 using Boc-pipecolic acid in place of Boc-thioprolin. Calculated weight 403.52, Found 371.1, 426.2(M-NHOH, M+Na).

20 **Synthesis of compound 89.** Synthesis of compound 89 was completed as described above for compound 64 using Boc-pipecolic acid in place of Boc-thioprolin. Calculated weight 396.51, Found 397.2, 419.0 (M+H, M+Na).

25 **Synthesis of compound 90.** Synthesis of compound 90 was completed as described above for compound 65 using Boc-pipecolic acid in place of Boc-thioprolin. Calculated weight 341.45, Found 364.2, (M+Na).

Synthesis of compound 91. Synthesis of compound 91 was completed as described above for compound 67 using Boc-pipecolic acid in place of Boc-thioprolin. Calculated weight 381.52, Found 349.2, 404.2 (M-NHOH, M+Na).

30 **Synthesis of compound 92.** Synthesis of compound 92 was completed as described above for compound 68 using Boc-pipecolic acid in place of Boc-thioprolin. Calculated weight 395.54, Found 363.2, 418.2 (M-NHOH, M+Na).

35

Synthesis of compound 93. Synthesis of compound 93 was completed as described above for compound 69 using Boc-pipecolic acid in place of Boc-thiopropine. Calculated weight 383.49, Found 406.2, (M+Na).


5 **Synthesis of compound 94.** Synthesis of compound 94 was completed as described above for compound 70 using Boc-pipecolic acid in place of Boc-thiopropine. Calculated weight 411.54, Found 434.2, (M+Na).

10 **Synthesis of compound 95.** Synthesis of compound 95 was completed as described above for compound 71 using Boc-pipecolic acid in place of Boc-thiopropine. Calculated weight 399.55, Found 367.3, 422.2, (M-NHOH, M+Na).

15 **Synthesis of compound 96.** Synthesis of compound 96 was completed as described above for compound 57 using Boc-pipecolic acid in place of Boc-thiopropine and N-methyl piperazine in place of isobutylamine. Calculated weight 396.53, Found 397.2, 420.2 (M+H, M+Na).

5.5.1 Synthesis of Compounds of Formula XVII

20 Compounds represented by Formula XVII are synthesized as described in section 5.5.1. See Table 13 for representative R groups.



Formula XVII

Synthesis of compound 97. Synthesis of compound 97 was completed as described above for compound using Boc-pipecolic acid in the place of Boc-thiopropine and pyrrolidine.

30 Step 1:

EDC (4 g, 20.86 mmol) and HOBT (4 g, 29.6 mmol) were added to a solution of Boc-pipecolic acid (4 g, 17.44 mmol) in DMF (60 ml) and stirred for 1 h. Isobutylamine (0.159 ml, 1.6 mmol) was added and the solution was stirred overnight. Saturated ammonium chloride solution was added (200 ml) and the mixture was extracted with
35 EtOAc. The organic layer was washed with saturated NaHCO₃, 0.5 NH₂Cl, brine and dried

over Na_2SO_4 . The solvent was removed in vacuo to give desired product. The Boc group was removed by treatment with 16 ml of 4.0 M HCl/dioxane for 2 h followed by removal of the solvent in vacuo to give the HCl salt of the amino amide. Crude Yield 3.4 g, quantitative.

5 Step 2:

This amino amide HCl salt (1.48 g, 6.76 mmol) was added along with triethylamine (0.139 ml, 1.00 mmol) to a solution of (R) tert-butyl (2-pentyl)succinate mono acid (1.5 g, 6.15 mmol), EDC (1.41 g, 7.38 mmol) and HOBT (1.40 g, 10.36 mmol) which had been stirred in 50 ml of DMF for 30 min. The resulting mixture was allowed to stir overnight. A
10 saturated solution of ammonium chloride (2.5 ml) was added and the mixture was extracted with EtOAc. The organic layer washed with saturated NaHCO_3 , 0.5 N HCl, brine and dried over Na_2SO_4 . The solvent was removed in vacuo to give crude oil which was purified by flash chromatography using 50% EtOAc in hexane. Yield 2.15g, 86%. Analyzed by LC/MS. Calculated weight 408.59; Found 338.3, 431.3 (M-C₄H₈N, M+Na).

15 Step 3:

The tert-butyl ester (1.6 g, 3.92 mmol) was subjected to 15 ml of 20% TFA in DCM and stirred for 1 h. Evaporation and dried under high vacuum to yield crude acid which was dissolved in 6 ml of methanol and 12 ml of benzene, and trimethylsilyldiazomethane in hexanes (2.0 M, 12.0 ml, 24 mmol) was added slowly. Volatile material was removed in
20 vacuo and the residue was purified by flash chromatography using EtOAc. Yield 1.05 g, 70%. Analyzed by LC/MS. Calculated weight 366.50; Found 296.3, 389.2 (M-C₄H₈N, M+Na).

Step 4:

A methanolic solution of NH_2OH HCl (2 M, 30 ml) was cooled to 0°C. A
25 methanolic solution of KOH (3 M, 30 ml) was added dropwise over 30 min, the mixture was stirred for 30 min and filtered. An aliquot (10 ml, 20 mmol NH_2OH) of this solution was added to the methyl ester (0.35 g, 0.95 mmol); product of the previous reaction. The mixture was allowed to shake for 30 min and acidified to pH 5 by Dowex acidic resin. The solvent was removed in vacuo and the residue was purified by flash chromatography using
30 5% MeOH in EtOAc. Yield 175 mg, 50%. Analyzed by LC/MS. Calculated weight 367.49; Found 297.2, 390.2 (M-C₄H₈N, M+Na).

Synthesis of compound 223 was completed as described above for compound 97 using D-Boc-pipecolic acid in place of L-Boc-pipecolic. Analyzed by HPLC/MS. Calculated
35 weight 367.49, Found 297.2, 368.3 (M-C₄H₈N, M+H).

Synthesis of compound 98. Synthesis of compound 98 was completed as described above for compound 97 using (R)-2-(t-butylcarboxymethyl)-3-(3,4-difluorophenyl)-propionic acid in place of (R) tert-butyl (2-pentyl) succinate mono acid. Analyzed by HPLC/MS.

Calculated weight 423.46, Found 446.1 (M+Na).

5

Synthesis of compound 99. Synthesis of compound 99 was completed as described above for compound 97 using (R)-2-(t-butylcarboxymethyl)-3-phenylmethylpropionic acid in place of (R) tert-butyl (2-pentyl) succinate mono acid. Analyzed by HPLC/MS. Calculated weight 387.48, Found 410.2 (M+Na).

10

Synthesis of compound 100. Synthesis of compound 100 was completed as described above for compound 97 using (R)-2-(t-butylcarboxymethyl)-3-cyclohexylmethylpropionic acid in place of (R) tert-butyl (2-pentyl) succinate mono acid. Analyzed by HPLC/MS. Calculated weight 393.53, Found 416.2 (M+Na).

15

Synthesis of compound 101. Synthesis of compound 101 was completed as described above for compound 97 using

(R)-2-(t-butylcarboxymethyl)-3-(3,4-difluorophenyl)-propionic acid in place of (R) tert-butyl (2-pentyl) succinate mono acid and 2-aminothiazole in place of pyrrolidine.

20 Analyzed by HPLC/MS. Calculated weight 452.48, Found 453.0, 475.0 (M+H, M+Na).

Synthesis of compound 102. Synthesis of compound 102 was completed as described above for compound 97 using (R)-2-(t-butylcarboxymethyl)-3-phenylmethylpropionic acid in place of (R) tert-butyl (2-pentyl) succinate mono acid and 2-aminothiazole in place of
25 pyrrolidine. Analyzed by HPLC/MS. Calculated weight 416.50, Found 417.1, 439.0 (M+H, M+Na).

Synthesis of compound 103. Synthesis of compound 103 was completed as described above for compound 97 using (R)-2-(t-butylcarboxymethyl)-3-cyclohexylmethylpropionic
30 acid in place of (R) tert-butyl (2-pentyl) succinate mono acid and 2-aminothiazole in place of pyrrolidine. Analyzed by HPLC/MS. Calculated weight 422.55, Found 423.1, 445.1 (M+H, M+Na).

35

Synthesis of compound 120AStep 1:

To a solution of (R) tert-butyl (2-pentyl)succinate mono N-hydroxysuccinimide ester (0.34 g, 1 mmol) in DCM was added N,N Diethylnipecotic acid (372 ml, 2.0 mmol) and TEA (2 ml, 14.65 mmol). The resulting mixture was stirred at room temperature for overnight and evaporated to dryness. The residue was taken up with EtOAc and extracted with 1 N HCl then brine. The organic layer was dried over anhydrous sodium sulfate, filtered and solvent removed under reduced pressure to give 0.48 g of crude product which was carried to next step without further purification.

10 Step 2:

Same as step 3 in the synthesis of compound 97


Step 3:

Same as step 4 in the synthesis of compound 97

15 5.2.6 Synthesis of Nipecotic Acid Series

Compounds represented by Formula XII were synthesized as described in section 5.2.2. See Scheme V for a general synthetic scheme and Table 9 for representative R groups. The nipecotic acid series was synthesized in an analogous fashion to the thioproline series (See 5.2.2 above), with Boc-thioproline replaced by Boc-nipecotic acid.

20



25

Formula XII

Synthesis of compound 104. Synthesis of compound 104 was completed as described above for compound 57 using Boc-nipecotic acid in place of Boc-thioproline.


30 Calculated weight 369.50, Found 337.2, 370.2, 392.3 (M-NHOH, M+H, M+Na).

Synthesis of compound 105. Synthesis of compound 105 was completed as described above for compound 58 using Boc-nipecotic acid in place of Boc-thioproline. Calculated weight 381.52, Found 349.2, 382.3, 404.2 (M-NHOH, M+H, M+Na).

35

Synthesis of compound 106. Synthesis of compound 106 as completed as described above for compound 59 using Boc-nipecotic acid in place of Boc-thiopropoline. Calculated weight 395.54, Found 363.3, 396.3, 418.0 (M-NHOH, M+H, M+Na).

5 **Synthesis of compound 107.** Synthesis of compound 107 was completed as described above for compound 60 using Boc-nipecotic acid in place of Boc-thiopropoline. Calculated weight 397.52, Found 365.2, 398.3, 420.2 (M-NHOH, M+H, M+Na).



Synthesis of compound 108. Synthesis of compound 108 was completed as described above for compound 61 using Boc-nipecotic acid in place of Boc-thiopropoline. Calculated weight 393.48, Found 361.2, 394.2, 416.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 109. Synthesis of compound 109 was completed as described above for compound 62 using Boc-nipecotic acid in place of Boc-thiopropoline. Calculated weight 409.55, Found 377.2, 410.2, 432.1 (M-NHOH, M+H, M+Na).

Synthesis of compound 110. Synthesis of compound 110 was completed as described above for compound 57 using Boc-nipecotic acid in place of Boc-thiopropoline and 2-aminomethylpyridine in place of isobutylamine. Calculated weight 404.51, Found 372.3, 405.2, 427.0 (M-NHOH, M+H, M+Na).

Synthesis of compound 111. Synthesis of compound 111 was completed as described above for compound 63 using Boc-nipecotic acid in place of Boc-thiopropoline. Calculated weight 403.52, Found 371.2, 404.2, 426.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 112. Synthesis of compound 112 was completed as described above for compound 64 using Boc-nipecotic acid in place of Boc-thiopropoline. Calculated weight 396.51, Found 364.1, 397.2, 419.0 (M-NHOH, M+H, M+Na).

Synthesis of compound 113. Synthesis of compound 113 was completed as described above for compound 65 using Boc-nipecotic acid in place of Boc-thioprolin. Calculated weight 341.45, Found 309.2, 342.2, 364.2 (M-NHOH, M+H, M+Na).

5 **Synthesis of compound 114.** Synthesis of compound 114 was completed as described above for compound 66 using Boc-nipecotic acid in place of Boc-thioprolin. Calculated weight 367.49, Found 335.2, 368.3, 390.0 (M-NHOH, M+H, M+Na).

10 **Synthesis of compound 115.** Synthesis of compound 115 was completed as described above for compound 67 using Boc-nipecotic acid in place of Boc-thioprolin. Calculated weight 381.52, Found 349.2, 382.3, 404.2 (M-NHOH, M+H, M+Na).

15 **Synthesis of compound 116.** Synthesis of compound 116 was completed as described above for compound 69 using Boc-nipecotic acid in place of Boc-thioprolin. Calculated weight 395.54, Found 363.3, 396.3, 418.2 (M-NHOH, M+H, M+Na).

20 **Synthesis of compound 117.** Synthesis of compound 117 was completed as described above for compound 70 using Boc-nipecotic acid in place of Boc-thioprolin. Calculated weight 383.49, Found 351.2, 384.2, 406.2 (M-NHOH, M+H, M+Na).

Synthesis of compound 118. Synthesis of compound 118 was completed as described above for compound 70 using Boc-nipecotic acid in place of Boc-thioprolin. Calculated weight 411.54, Found 379.2, 412.3, 434.2 (M-NHOH, M+H, M+Na).


25 **Synthesis of compound 119.** Synthesis of compound 119 was completed as described above for compound 71 using Boc-nipecotic acid in place of Boc-thioprolin. Calculated weight 399.55, Found 367.3, 400.2, 422.1 (M-NHOH, M+H, M+Na).

30 **Synthesis of compound 120.** Synthesis of compound 120 was completed as described above for compound 57 using Boc-nipecotic acid in place of Boc-thioprolin and N-methyl piperazine in place of isobutylamine. Calculated weight 396.53, Found 364.2, 397.2 (M-NHOH, M+H).

35

5.2.7 Synthesis of Isonipectic Acid Series


Compounds represented by Formula XIII are synthesized as described in section 5.2.2. See Scheme VI for a general synthetic scheme and Table 10 for representative R groups. The isonipecotic acid series was synthesized in an analogous fashion to the thioproline series, wherein Boc-thioproline replaced with Boc-isonipacotic acid.



Formula XIII

Synthesis of compound 123. Synthesis of compound 123 was completed as described for compound 59 above for using Boc-isonipectic acid in place of Boc-thioproline. Calculated weight 369.51, Found 337.2, 370.2, 392.3 (M-NHOH, M+H, M+Na).

Synthesis of compound 124. Synthesis of compound 124 was completed as described for compound 58 above for using Boc-isonipectic acid in place of Boc-thioproline. Calculated weight 381.52, Found 349.2, 382.3 (M-NHOH, M+H).



Scheme VI

Synthesis of compound 125. Synthesis of compound 125 was completed as described for compound 61 above for using Boc-isonipecotic acid in place of Boc-thioprolin. Calculated weight 393.49, Found 361.2, 394.2 (M-NHOH, M+H).

5 **Synthesis of compound 126.** Synthesis of compound 126 was completed as described for compound 62 above for using Boc-isonipecotic acid in place of Boc-thioprolin. Calculated weight 409.55, Found 377.2, 410.2 (M-NHOH, M+H).

10 **Synthesis of compound 127.** Synthesis of compound 127 was completed as described for compound 57 above for using Boc-isonipecotic acid in place of Boc-thioprolin and 2-aminomethylpyridine in place of isobutylamine. Calculated weight 404.51, Found 372.3, 405.2 (M-NHOH, M+H).

15 **Synthesis of compound 128.** Synthesis of compound 128 was completed as described for compound 63 above for using Boc-isonipecotic acid in place of Boc-thioprolin. Calculated weight 403.53, Found 371.2, 404.2, 426.2 (M-NHOH, M+H, M+Na).

20 **Synthesis of compound 129.** Synthesis of compound 129 was completed as described for compound 64 above for using Boc-isonipecotic acid in place of Boc-thioprolin. Calculated weight 369.51, Found 364.2, 397.2 (M-NHOH, M+H).

25 **Synthesis of compound 130.** Synthesis of compound 130 was completed as described for compound 65 above for using Boc-isonipecotic acid in place of Boc-thioprolin. Calculated weight 341.45, Found 309.2, 342.3, 364.2 (M-NHOH, M+H, M+Na).

30 **Synthesis of compound 131.** Synthesis of compound 131 was completed as described for compound 66 above for using Boc-isonipecotic acid in place of Boc-thioprolin. Calculated weight 367.49, Found 335.2, 368.3, 390.2 (M-NHOH, M+H, M+Na).

35 **Synthesis of compound 132.** Synthesis of compound 132 was completed as described for compound 67 above for using Boc-isonipecotic acid in place of Boc-thioprolin. Calculated weight 381.52, Found 349.2, 382.3 (M-NHOH, M+H).

Synthesis of compound 133. Synthesis of compound 133 was completed as described for compound 68 above for using Boc-isonipecotic acid in place of Boc-thiopropine. Calculated weight 395.55, Found 363.3, 396.3 (M-NHOH, M+H).

5 **Synthesis of compound 134.** Synthesis of compound 134 was completed as described for compound 69 above for using Boc-isonipecotic acid in place of Boc-thiopropine. Calculated weight 383.49, Found 351.2, 384.2 (M-NHOH, M+H).

10 **Synthesis of compound 135.** Synthesis of compound 135 was completed as described for compound 70 above for using Boc-isonipecotic acid in place of Boc-thiopropine. Calculated weight 411.55, Found 379.2, 412.3 (M-NHOH, M+H).

15 **Synthesis of compound 136.** Synthesis of compound 136 was completed as described for compound 71 above for using Boc-isonipacotic acid in place of Boc-thiopropine. Calculated weight 399.56, Found 367.3, 400.2 (M-NHOH, M+H).

5.5.8 Aminoalcohol Series

Synthesis of aminoalcohol series of compounds was completed as described below using the following 7 amines: 2-piperidinemethanol, 4-piperidineethanol, 20 N-benzylethanolamine, diethanolamine, N-methylhomoveratrylamine, 2-(methylamino)ethanol, 3-aminobenzyl alcohol, 4-aminophenylethyl alcohol. See Scheme VII below and Table 11 for representative R groups.



30 Scheme VII

Step 1:

Amines (0.18 mmol) were added to each vial in a 4 x 6 array of tubes. To each was added 0.6 ml of DMF, DIEA (0.091 ml, 0.525 mmol), and chlorotrimethylsilane (0.040 ml, 3.15 mmol), and the array was shaken for 2.5 h.

35

To a separate solution of (R) methyl (2-pentyl)succinate mono acid (1.0 g, 5.0 mmol) in DMF (14 ml) at 0°C was added EDC (0.96 g, 5.0 mmol) and DIEA (209 µl, 1.2 mmol). This mixture was stirred for 20 min, and aliquots (0.42 ml, 0.15 mmol activated acid) were added to each vial in the amine array, followed by a catalytic amount (ca. 5 mg) of DMAP. The mixtures were shaken for 18 h, diluted with 3 ml ethyl acetate, washed with 10% NaHSO₄ (aq) and brine, filtered through Na₂SO₄ and evaporated. The resulting methyl esters were placed in a vacuum chamber under high vacuum overnight, weighed, and analyzed by HPLC/MS.

Step 2:

A methanolic solution of NH₂OH HCl (2 M, 15 ml) was cooled to 0°C. A methanolic solution of KOH (3 M, 15 ml) was added and the mixture was stirred for 30 min and filtered. An aliquot (2 ml, 2.0 mmol NH₂OH) of this solution was added to each of the reaction vials containing the methyl ester product of the previous reaction. The array was shaken for 30 min and Dowex acidic resin (220 mg, 1.125 mmol) was added to each vial and the solutions were filtered, evaporated and purified by preparative HPLC. The product-containing fractions were collected, evaporated in a Savant evaporating centrifuge, and re-analyzed by HPLC/MS.

Synthesis of compound 201: Calculated weight 330.40, Found 268.2, 301.2, 323.2 (M-NHOH, M+H, M+Na)

Synthesis of compound 202: Calculated weight 314.43, Found 282.3, 315.2, 338.2 (M-NHOH, M+H, M+Na)

Synthesis of compound 203: Calculated weight 336.43, Found 304.2, 337.1, 359.0 (M-NHOH, M+H, M+Na)

Synthesis of compound 204: Calculated weight 290.36, Found 258.1, 291.3, 313.2 (M-NHOH, M+H, M+Na)

Synthesis of compound 210: Calculated weight 380.48, Found 348.3, 381.3, 403.2 (M-NHOH, M+H, M+Na)

Synthesis of compound 211: Calculated weight 260.33, Found 228.2, 261.2, 283.2 (M-NHOH, M+H, M+Na)

Synthesis of compound 212: Calculated weight 308.38, Found 276.1, 309.2, 331.1
(M-NH₂OH, M+H, M+Na)

Synthesis of compound 213: Calculated weight 322.40, Found 290.3, 323.3, 345.1
5 (M-NH₂OH, M+H, M+Na)

Piperazine Series I

See Scheme VIII below. See Table 11 for representative R groups.

Step 1:

10 **Synthesis of compound 217.** To a stirred solution of piperazine (6.1 g, 71 mmol) in CH₂Cl₂ (50 ml) was added a solution of benzoyl chloride (1.0 g, 7 mmol) in CH₂Cl₂ (50 ml) at 0°C. The reaction mixture was stirred overnight. After addition of H₂O, the mixture was extracted with CHCl₃, washed with H₂O, then dried (MgSO₄) and concentrated. Purification of the residue by chromatography eluting with CHCl₃ / MeOH gave benzoyl
15 piperazine (393 mg).

Step 2 :

To a mixture of (R)-tert-butyl (2-pentyl)succinate mono N-hydroxysuccinimide ester (500mg, 1.5mmol), benzoylpiperazine (380mg, 2.0 mmol) in CH₂Cl₂ (5.3 ml) was added Et₃N (278 ml, 2.0 mmol) at 0°C. The resulting mixture was stirred at room temperature for
20 2.5 days, then concentrated and diluted with EtOAc. The organic phase was successively washed with 1N HCl, saturated NaHCO₃, brine, and then dried (MgSO₄) and concentrated. Purification of the residue by chromatography eluting with n-Hexane / EtOAc gave the 4-((R)-tert-butyl (2-pentyl)succinate) benzoyl piperazine (383 mg).


Step 3:

25 The 4-((R)-tert-butyl (2-pentyl)succinate) benzoyl piperazine (378 mg, 1.0 mmol) was dissolved in CH₂Cl₂ (4 ml) and trifluoroacetic acid (1.6 ml) was added at 0°C. The reaction mixture was stirred at room temperature for 3 h and concentrated. After addition of H₂O, the mixture was extracted with CH₂Cl₂, dried (MgSO₄) and concentrated to give the acid (193mg).

Step 4:

30 To a stirring solution of the acid (190 mg, 0.53 mmol), and Et₃N (96 ml, 0.69 mmol) in THF (8 ml) was added isobutyl chloroformate (82 ml, 0.63 mmol) at 0°C. The mixture was stirred at 0°C for 15 min. TMSONH₂ was then added and allowed to stir for 2 h. EtOAc was added to the solution and washed with saturated NH₄Cl. The mixture was dried
35 (MgSO₄) and concentrated. Purification of the residue was accomplished by

chromatography with the eluants EtOAc then CHCl₃/MeOH (10:1) and gave compound 217 (97 mg). Analyzed by ¹H NMR (DMSO-d₆) δ: 10.46 (s, 1H), 8.77 (s, 1H), 7.6-7.45 (m, 5H), 3.9-3.1 (m, 9H), 2.34 (dd, J=14.4Hz, 8.4 Hz, 1H), 2.12 (dd, J=15.0Hz, 6.0Hz, 1H), 1.63-1.2 (m, 8H), 1.0-0.85 (m, 3H). HPLC/MS. Calculated weight 375.47, Found 374.2 (M-H).



Scheme VIII

20 Synthesis of compound 218.

Piperazine series II

See Scheme IX. See table 11 for representative R groups.

Step 1:


To a stirred solution of Boc-piperazine (1.5 g, 8.1 mmol) and Et₃N (0.9 g, 8.9 mmol) in CH₂Cl₂ (20 ml) was added 1-pyrrolidinecarbonyl chloride (0.89 ml, 8.1 mmol) at 0°C and stirred at room temperature for 3 days. After addition of H₂O, the mixture was extracted with CH₂Cl₂, washed with H₂O, then dried (MgSO₄) and concentrated. Purification of the residue was accomplished by chromatography with the eluant EtOAc/n-Hexane (1:10 to 1:2) to give the Boc protected urea (2.18 g).

Step 2:

The Boc protected urea (1.0 g, 3.5 mmol) was dissolved in CH₂Cl₂ (10 ml) and TFA (4 ml) was added at 0°C. The reaction mixture was stirred overnight at room temperature and concentrated. 10% NaOH was added and the pH of the mixture was brought to 10. The mixture was extracted with CHCl₃, dried (MgSO₄) and concentrated to give the urea (193 mg).

Steps 3-5:

Steps 3-5 in the synthesis of compound 218 were completed as described above in Step 2-4 for the synthesis of compound 217 using the urea from step 2 in place of N-benzoylpiperazine. Analyzed by ^1H NMR (DMSO- d_6) δ : 10.45 (s, 1H), 8.77 (s, 1H), 4.1-2.7 (m, 13H), 2.32 (dd, $J=14.7$ Hz, 8.1 Hz, 1H), 2.10 (dd, $J=14.7$ Hz, 5.9 Hz, 1H), 1.9-1.75 (m, 4H), 1.6-1.1 (m, 8H), 0.97-0.85 (m, 3H). HPLC/MS. Calculated weight 368.48. Found 336.4, 369.4, 391.2 (M-NHOH, M+H, M+Na).



Scheme IX

20 Aminopyrrolidine

See Scheme X. See Table 11 for representative R groups.

Step 1:

Step 1 was completed as described above in Step 1 for the synthesis of compound 217 using (3S)-(+)-1-benzyl-3-amino pyrrolidine in place of piperazine.


25 Step 2:

(3S)-1-benzyl-3-(N-benzoylamino) pyrrolidine (400 mg, 1.4 mmol) was dissolved in MeOH (6 ml) and 10% Pd/C (400 mg) was added at 0°C. After addition of HCO_2NH_4 (450 mg, 7.1 mmol), the reaction mixture was heated under reflux for 10 min. The catalyst was filtered off and the filtrate was concentrated to give (3S)-3-(N-benzoylamino) pyrrolidine (180 mg). MS (ES) 191.3 (M+H)

Steps 3-5:

Steps 3-5 in the synthesis of compound 219 were completed as described above in Step 2-4 for the synthesis of compound 217 using (3S)-3-(N-benzoylamino) pyrrolidine in place of N-benzoylpiperazine. Analyzed by ^1H NMR (DMSO- d_6) δ : 10.55-10.45 (m, 1H), 8.8 (s, 1H), 8.65-8.5 (m, 1H), 8.0-7.5 (m, 5H), 4.6-4.45 (m, 1H), 4.0-3.8 (m, 1H), 3.8-3.35

(m, 3H), 3.05-2.8 (m, 1H), 2.4-1.8 (m, 5H), 1.6-1.0 (m, 8H), 0.95-0.75 (m, 3H). HPLC/MS
Calculated weight 375.47, Found 343.4, 376.3, 398.2 (M-NHOH, M+H, M+Na).



Scheme X

Hydroxyazetidine Series

See Scheme XI. See Table 11 for representative R groups.

Step 1:

15 Synthesis of compound 221. Step 1 was completed as described above in Step 2 of the synthesis of compound 219 using 1-diphenyl-3-hydroxyazetidine in place of (3S)-1-benzyl-3-(N-benzoylamino) pyrrolidine.

Step 2:

20 Step 2 was completed as described above in Step 3 of the synthesis of compound 218 using 3-hydroxyazetidine in place of benzoylpiperazine.

Step 3:

25 A mixture of the alcohol formed in Step 2 (165 mg, 0.55 mmol), phenyl isocyanate (72 ml, 0.66 mmol), Et₃N (154 ml, 1.10 mmol), DMAP (8 mg, 0.07 mmol) in THF (8 ml) was stirred at 70°C for overnight. EtOAc was added and washed with 1N HCl, saturated Na₂CO₃, brine. The organic layer was dried (MgSO₄) and concentrated, followed by addition of PhCH₃. Precipitate was filtered off and filtrate was concentrated to give crude (R)-tert-butyl (2-pentyl)succinate amide (250 mg), which was used without further purification.


Steps 4-5:

30 Steps 4-5 in the synthesis of compound 221 were completed as described above in Step 3-4 for the synthesis of compound 218 using the (R)-tert-butyl (2-pentyl)succinate amide in place of N-benzoylpiperazine. Analyzed by ¹H NMR (DMSO-d₆) δ: 10.45, 10.43 (2s, 1H), 9.96 (s, 1H), 8.8, 8.76 (2s, 1H), 7.6-7.4 (m, 2H), 7.4-7.25 (m, 2H), 7.0-7.1 (m, 1H), 5.3-5.1 (m, 1H), 4.7-4.45 (m, 1H), 4.35-4.0 (m, 2H), 3.9-3.7 (m, 1H), 2.75-2.6

35

(m, 1H), 2.33-2.13 (m, 1H), 2.13-1.97 (m, 1H), 1.55-0.95 (m, 8H), 0.95-0.8 (m, 3H).

HPLC/ MS. Calculated weight 377.44. Found 345.3, 378.3 (M-NHOH, M+H).




Scheme XI

15 Aminoazetidine

Synthesis of compound 222 was completed as described above for compound 219 using 1 diphenylmethyl-3-amino azetidine in place of (3S)-(+)-1-benzyl-3-amino pyrrolidine. See Scheme XII. Analyzed by ¹H NMR (DMSO-d₆) δ: 9.1 (br s, 1H), 8.1-7.8 (m, 2H), 7.7-7.4 (m, 3H), 4.85-3.0 (m, 4H), 2.9-1.9 (m, 4H), 1.6-1.0 (m, 8H), 1.0-0.75 (m, 3H). HPLC/ MS. Calculated weight 361.44. Found 329.2, 362.3 (M-NHOH, M+H).

20



Scheme XII

35

5.6.1 Synthesis of Related Deformylase Inhibitors

Compounds represented by the Formulas I and II are synthesized as described in section 5.6.1. See Scheme XIII for a general synthetic scheme and Table 11 for representative R groups. Boc-protected amino acids from the group comprising Val, Nva, Phe, Met, Ser(OBn), and Tyr(OBn) were individually treated with amine reagents in the presence of EDC to afford Boc-protected amino amides. Deprotection of the Boc group followed by addition of (R)-tert-butyl (2-pentyl)succinate mono acid and EDC afforded tert-butyl ester substituted amides. Conversion of the tert-butyl esters to the carboxylic acids with subsequent treatment with trimethylsilyldiazomethane afforded the methyl ester. Finally, addition of hydroxylamine hydrochloride yielded the desired products.

Step 1:

Six Boc-protected amino acids (Val, Nva, Phe, Met, Ser(OBn), Tyr(OBn)) were each dissolved in DCM to give a concentration of 1 M, and 4 aliquots (0.48 ml, 0.48 mmol) of each were added to each column of the array, one per vial. Three of the starting amine reagents (pyrrolidine, piperidine, and benzylamine) were each dissolved in DCM at 2.0 M. Dimethylamine was used as provided, in a 2 M THF solution (Aldrich). Six aliquots (0.30 ml, 0.6 mmol) of each of the 4 amine solutions were added to each row of the array, one per vial. A 0.25 M solution of EDC in DCM was prepared and an aliquot (2.40 ml, 0.6 mmol) was added to each vial. The vials were sealed in the block and agitated at 200 rpm on a rotary shaker for 18 h. To each vial was added 1.5 ml 10% NaHSO₄ (aq), the array was sealed with the membrane, agitated, and the aqueous layers were withdrawn. The wash procedure was repeated with 10% NaHSO₄, NaHCO₃ (saturated), and brine. The solutions were filtered through approximately 0.5 g anhydrous Na₂SO₄ each and evaporated to dryness. The resulting Boc-protected amides (oils and solids) were placed in a vacuum chamber under high vacuum overnight, weighed, and analyzed by HPLC/MS.

Step 2:

Deprotection of the Boc-protected amino amides was accomplished by dissolving each product in 2 ml of 4 M HCl in dioxane, followed by shaking for 45 min, and evaporation of solvents. The vials were then placed under high vacuum overnight, weighed, and the amine hydrochlorides were analyzed by HPLC/MS.

Step 3:

To each of the amine hydrochlorides was added a solution of (R) tert-butyl (2-pentyl)succinate mono acid in DCM (2.0 M, 0.20 ml, 0.40 mmol), EDC in DCM (0.25 M, 2.4 ml, 0.60 mmol), and DIEA (209 ml, 1.2 mmol). The mixtures were agitated for 18 h, washed twice with 10% NaHSO₄ (aqueous), once each with NaHCO₃ (saturated), and

brine, filtered through Na_2SO_4 and evaporated. The resulting tert-butyl ester substituted amides were then weighed and analyzed by HPLC/MS.

Step 4:

5 The tert-butyl esters were converted to the carboxylic acids by dissolving each in 2 ml of 4 M HCl in dioxane, shaken for 45 min, and evaporated. The vials were then placed under high vacuum overnight, weighed, and the acids were analyzed by HPLC/MS.

Step 5:

To convert the acids to the corresponding methyl esters each was dissolved in 0.3 ml of methanol and 0.7 ml of benzene, and trimethylsilyldiazomethane in hexanes (2.0 M, 0.60 10 ml, 1.2 mmol) was added slowly. The array was agitated for 2 h, the solutions were evaporated, and the residues placed under high vacuum overnight. The resulting methyl esters were weighed and analyzed by HPLC/MS.


Step 6:

A methanolic solution of $\text{NH}_2\text{OH HCl}$ (2 M, 30 ml) was cooled to 0°C . A 15 methanolic solution of KOH (3 M, 30 ml) was added dropwise over 30 min, the mixture was stirred for 30 min and filtered. An aliquot (2 ml, 2.0 mmol NH_2OH) of this solution was added to each of the reaction vials containing the methyl ester product of the previous reaction. The array was agitated for 30 min. Each solution was partially neutralized with 0.10 ml of conc. HCl, then quickly buffered with 0.30 ml of 1 M triethylammonium 20 bicarbonate (pH 7). The solutions were then filtered and directly purified by preparative HPLC. The product-containing fractions were collected, evaporated in a Savant evaporating centrifuge, and re-analyzed by HPLC/MS.


25

30

35



5



Scheme XIII

Synthesis of compound 137 was accomplished as described in the Section 5.6 calculated
 15 M.W. 355.48 found M.W. (M-NHOH, M+H, M+Na) 323.1, 356.3, 378.2.

Synthesis of compound 138 was accomplished as described in the Section 5.6 calculated
 M.W. 329.44 found M.W. (M-NHOH, M+H, M+Na) 297.1, 330.2, 352.2.

20 Synthesis of compound 139 was accomplished as described in the Section 5.6 calculated
 M.W. 377.48 found M.W. (M-NHOH, M+H, M+Na) 345.1, 378.2, 400.1.

Synthesis of compound 140 was accomplished as described in the Section 5.6 calculated
 M.W. 361.51 found M.W. (M-NHOH, M+H, M+Na) 329.1, 362.1, 384.1.

25

Synthesis of compound 141 was accomplished as described in the Section 5.6 calculated
 M.W. 407.51 found M.W. (M-NHOH, M+H, M+Na) 375.2, 408.3, 430.2.

Synthesis of compound 142 was accomplished as described in the Section 5.6 calculated
 30 M.W. 483.61 found M.W. (M-NHOH, M+H, M+Na) 451.2, 484.2, 506.2.

Synthesis of compound 143 was accomplished as described in the Section 5.6 calculated
 M.W. 355.48 found M.W. (M-NHOH, M+H, M+Na) 323.1, 356.3, 378.3.

35

Synthesis of compound 144 was accomplished as described in the Section 5.6 calculated M.W. 403.52 found M.W. (M-NHOH, M+H, M+Na) 371.2, 404.2, 426.2.

5 Synthesis of compound 145 was accomplished as described in the Section 5.6 calculated M.W. 387.54 found M.W. (M-NHOH, M+H, M+Na) 355.1, 388.1, 410.1.

Synthesis of compound 146 was accomplished as described in the Section 5.6 calculated M.W. 433.55 found M.W. (M-NHOH, M+H, M+Na) 401.2, 434.2, 456.1.

10 Synthesis of compound 149 was accomplished as described in the Section 5.6 calculated M.W. 369.50 found M.W. (M-NHOH, M+H, M+Na) 337.2, 370.2, 393.2.

Synthesis of compound 150 was accomplished as described in the Section 5.6 calculated M.W. 417.55 found M.W. (M-NHOH, M+H, M+Na) 385.2, 418.2, 440.2.

15

Synthesis of compound 151 was accomplished as described in the Section 5.6 calculated M.W. 401.57 found M.W. (M-NHOH, M+H, M+Na) 369.1, 402.2, 424.2.

20 Synthesis of compound 152 was accomplished as described in the Section 5.6 calculated M.W. 447.58 found M.W. (M-NHOH, M+H, M+Na) 415.3, 448.23, 470.2.

Synthesis of compound 153 was accomplished as described in the Section 5.6 calculated M.W. 523.67 found M.W. (M-NHOH, M+H, M+Na) 491.2, 524.3, 546.3.

25 Synthesis of compound 154 was accomplished as described in the Section 5.6 calculated M.W. 391.51 found M.W. (M-NHOH, M+H, M+Na) 359.2, 392.3, 414.2.

Synthesis of compound 155 was accomplished as described in the Section 5.6 calculated M.W. 391.51 found M.W. (M-NHOH, M+H, M+Na) 359.2, 392.3, 414.2.

30

Synthesis of compound 156 was accomplished as described in the Section 5.6 calculated M.W. 439.55 found M.W. (M-NHOH, M+H, M+Na) 407.3, 440.2, 462.2.

35 Synthesis of compound 157 was accomplished as described in the Section 5.6 calculated M.W. 423.58 found M.W. (M-NHOH, M+H, M+Na) 391.1, 424.1, 446.1.

Synthesis of compound 158 was accomplished as described in the Section 5.6 calculated M.W. 469.58 found M.W. (M-NH₂OH, M+H, M+Na) 437.2, 470.2, 492.3.

5 Synthesis of compound 159 was accomplished as described in the Section 5.6 calculated M.W. 545.68 found M.W. (M-NH₂OH, M+H, M+Na) 513.3, 546.3, 568.3.

Synthesis of compound 161 was accomplished as described in the Section 5.6 calculated M.W. 258.36 found M.W. (M-NH₂OH, M+H, M+Na) 226.36, 258.2, 281.4.

10 Synthesis of compound 162 was accomplished as described in the Section 5.6 calculated M.W. 272.39 found M.W. (M-NH₂OH, M+H, M+Na) 240.4, 273.4, 295.4.

Synthesis of compound 163 was accomplished as described in the Section 5.6 calculated M.W. 256.35 found M.W. (M-NH₂OH, M+H, M+Na) 224.4, 257.4, 279.2.

15 Synthesis of compound 164 was accomplished as described in the Section 5.6 calculated M.W. 270.37 found M.W. (M-NH₂OH, M+H, M+Na) 238.4, 271.4, 293.4.

20 Synthesis of compound 165 was accomplished as described in the Section 5.6 calculated M.W. 282.34 found M.W. (M-NH₂OH, M+H, M+Na) 250.3, 283.2, 305.2.

Synthesis of compound 166 was accomplished as described in the Section 5.6 calculated M.W. 336.47 found M.W. (M-NH₂OH, M+H, M+Na) 304.4, 337.4, 359.4.

25 Synthesis of compound 167 was accomplished as described in the Section 5.6 calculated M.W. 278.35 found M.W. (M-NH₂OH, M+H, M+Na) 246.3, 279.2, 301.2.

Synthesis of compound 168 was accomplished as described in the Section 5.6 calculated M.W. 363.46 found M.W. (M-NH₂OH, M+H, M+Na) 331.3, 364.4, 386.3.

30 Synthesis of compound 169 was accomplished as described in the Section 5.6 calculated M.W. 364.49 found M.W. (M-NH₂OH, M+H, M+Na) 332.4, 365.3, 387.2.

35 Synthesis of compound 170 was accomplished as described in the Section 5.6 calculated M.W. 296.34 found M.W. (M-NH₂OH, M+H, M+Na) 264.2, 297.2, 319.2

Synthesis of compound 171 was accomplished as described in the Section 5.6 calculated M.W. 357.25 found M.W. (M-NHOH, M+H, M+Na) 381.0, (Br isotopes) 324.2, 326.2, 357.2, 359.2, 379.0, 381.0.

- 5 Synthesis of compound 172 was accomplished as described in the Section 5.6 calculated M.W. 357.25 found M.W. (M-NHOH, M+H, M+Na) (Br isotopes) 324.2, 326.2, 357.2, 359.2, 379.0, 381.0.

- Synthesis of compound 173 was accomplished as described in the Section 5.6 calculated
10 M.W. 312.80 found M.W. (M-NHOH, M+H, M+Na) (Cl isotopes) 280.2, 282.1, 313.3, 315.2, 335.2, 337.2.

- Synthesis of compound 174 was accomplished as described in the Section 5.6 calculated M.W. 338.40 found M.W. (M-NHOH, M+H, M+Na) 306.2, 339.3, 361.2.

- 15 Synthesis of compound 173 was accomplished as described in the Section 5.6 calculated M.W. 370.45 found M.W. (M-NHOH, M+H, M+Na) 338.3, 371.2, 393.2.

- Synthesis of compound 176 was accomplished as described in the Section 5.6 calculated
20 M.W. 370.45 found M.W. (M-NHOH, M+H, M+Na) 338.2, 371.3, 393.3.

Synthesis of compound 177 was accomplished as described in the Section 5.6 calculated M.W. 335.40 found M.W. (M-NHOH, M+H, M+Na) 303.2, 336.2, 358.2.

- 25 Synthesis of compound 183 was accomplished as described in the Section 5.6 calculated M.W. 398.55 found M.W. (M-NHOH, M+H, M+Na) 399.4

- Synthesis of compound 184 was accomplished as described in the Section 5.6 calculated M.W. 349.43 found M.W. (M-NHOH, M+H, M+Na) 317.4, 350.4, 372.2.

- 30 Synthesis of compound 185 was accomplished as described in the Section 5.6 calculated M.W. 420.55 found M.W. (M-NHOH, M+H, M+Na) 421.4.

- Synthesis of compound 186 was accomplished as described in the Section 5.6 calculated
35 M.W. 365.43 found M.W. (M-NHOH, M+H, M+Na) 333.3, 366.4.

Synthesis of compound 187 was accomplished as described in the Section 5.6 calculated M.W. 436.55 found M.W. (M-NHOH, M+H, M+Na) 437.3.

5 Synthesis of compound 188 was accomplished as described in the Section 5.6 calculated M.W. 438.48 found M.W. (M-NHOH, M+H, M+Na) 406.3, 439.2, 461.22.

Synthesis of compound 189 was accomplished as described in the Section 5.6 calculated M.W. 452.51 found M.W. (M-NHOH, M+H, M+Na) 420.2, 453.3.

10 Synthesis of compound 190 was accomplished as described in the Section 5.6 calculated M.W. 353.39 found M.W. (M-NHOH, M+H, M+Na) 321.12, 353.4.

Synthesis of compound 191 was accomplished as described in the Section 5.6 calculated M.W. 367.42 found M.W. (M-NHOH, M+H, M+Na) 3335.3, 368.4, 390.2.

15

Synthesis of compound 182 was accomplished as described in the Section 5.6 calculated M.W. 358.48 found M.W. (M-NHOH, M+H, M+Na) 359.3.

20 Synthesis of compound 178 was accomplished as described in the Section 5.6 calculated M.W. 317.39 found M.W. (M-NHOH, M+H, M+Na) 285.4, 318.3, 340.4.

Synthesis of compound 179 was accomplished as described in the Section 5.6 calculated M.W. 384.52 found M.W. (M-NHOH, M+H, M+Na) 385.4.

25 Synthesis of compound 180 was accomplished as described in the Section 5.6 calculated M.W. 386.45 found M.W. (M-NHOH, M+H, M+Na) 354.4, 387.4, 409.3.

Synthesis of compound 181 was accomplished as described in the Section 5.6 calculated M.W. 327.42 found M.W. (M-NHOH, M+H, M+Na) 295.4, 328.4, 350.3.

30

5.6.2 Synthesis of Compounds of Formulas I and II

Compounds represented by Formulas I and II are synthesized as described in section 5.6.2. See Scheme XIV for a general synthetic scheme and Table 11 for representative R groups.

35

The following series of 24 amines was used: butylamine, amylamine, diethylamine, pyrrolidine, piperidine, morpholine, furfurylamine, Boc-piperizine, 1-adamantylamine, aniline, 4-morpholinoaniline, 4-pentyloxyaniline, 2-fluoroaniline, 3-fluoroaniline, 4-fluoroaniline, 3-bromoaniline, 4-bromoaniline, 4-chloroaniline, 3,5-dimethoxyaniline, 3-phenoxyaniline, 4-phenoxyaniline, 3-aminobenzamide, 4-aminobenzamide, 4-aminoacetanilide.

Step 1:

The series of 24 amines was added to each vial in a 4 x 6 array of tubes. To each was added 1 ml of DCM and a solution of (R) tert-butyl (2-pentyl)succinate mono N-hydroxysuccinimide ester in DCM (0.25 M, 1.20 ml, 0.30 mmol). The mixtures were agitated for 18 h, diluted to 2.5 ml with DCM, washed twice with 10% NaHSO₄ (aqueous), once each with NaHCO₃ (saturated), and brine, filtered through Na₂SO₄ and evaporated. The resulting tert-butyl ester substituted amides were then weighed and analyzed by HPLC/MS.


Small amounts of unreacted succinate-NHS ester from the mixtures were removed by diluting with 1 ml of DCM and approximately 50 mg of tris(2-aminoethyl)amine, polymer-bound (4.4 mmol N/g, Aldrich) was added. The mixtures were agitated for 18 h, filtered, evaporated, and analyzed by HPLC/MS.

Step 2:

The tert-butyl esters were converted to the carboxylic acids by dissolving each in 2 ml of a solution of 100:5:1 TFA:water:DMS, followed by shaking for 45 min, and evaporation of solvents. The vials were then placed under high vacuum overnight, weighed, and the acids were analyzed by HPLC/MS.

Steps 3-4:

Performed as in steps 5 and 6 in Section 5.6.1, as described above.



Scheme XIV

35

5.6.3 Synthesis of Compounds of Formulas I and II

Compounds represented by Formulas I and II are synthesized as described in section 5.6.3. See Scheme XIV for a general synthetic scheme and Table 11 for representative R groups.

- 5 These compounds were synthesized in a procedure similar to the above series, except that 0.36 mmol of Boc-protected amino acid starting materials were used and all other reagents were scaled appropriately. Also, N-hydroxysuccinimide (NHS) esters were used as coupling partners in Step 1 and Step 3 instead of the free acids and EDC.

Step 1:

- 10 Six Boc-protected amino acid NHS esters (Gly, b-Ala, Lys(Z), Ser(OBn), Asp(OBn, Glu(OBn)) were each dissolved in DCM to a concentration of 0.75 M, and 4 aliquots (0.48 ml, 0.36 mmol) of each were added to each column of the array, one per vial. Three of the starting amine reagents (pyrrolidine, piperidine, and benzylamine) were each dissolved in DCM at 2.0 M. Dimethylamine was used as provided, in a 2 M THF solution (Aldrich).
- 15 Six aliquots (0.225 ml, 0.45 mmol) of each of the 4 amine solutions were added to each row of the array, one per vial. The mixtures were agitated for 18 h, washed twice with 10% NaHSO₄ (aqueous), once each with NaHCO₃ (saturated), and brine, filtered through Na₂SO₄ and evaporated. The resulting Boc-protected amides (oils and solids) were placed in a vacuum chamber under high vacuum overnight, weighed, and analyzed by HPLC/MS.

20 Step 2:

Performed as in Step 2 of Section 5.6.2, as described above.

Step 3:


- To each of the amine hydrochlorides from Step 2 was added a solution of (R) tert-butyl (2-pentyl)succinate mono N-hydroxysuccinimide ester in DCM (1.5 M, 0.20 ml, 25 0.30 mmol), and DIEA (209 ml, 1.2 mmol). The mixtures were agitated for 18 h, washed twice with 10% NaHSO₄ (aq), once each with NaHCO₃ (saturated), and brine, filtered through Na₂SO₄ and evaporated. The resulting tert-butyl ester substituted amides were then weighed and analyzed by HPLC/MS.

30

35

Steps 4-6:

Performed as in steps 4-6 of Section 5.6.2, as described above.



Scheme XV

15 5.6.4 Synthesis of Compounds of Formulas I and II

Compounds represented by the Formulas I and II are synthesized as described in section 5.6.4. See Scheme XIII for a general synthetic scheme and Table 11 for representative R groups.

20 These compounds were synthesized in a procedure similar to the above series, except that the amine reagents used were aniline, p-toluidine, p-anisidine, and p-fluoroaniline.

The group of 24 amines used was: 3-hydroxypiperidine hydrochloride, 4-hydroxypiperidine, 2-piperidinemethanol, 3-piperidinemethanol, 2-piperidineethanol, 4-piperidineethanol, N-benzylethanolamine, diethanolamine, 1-(2-hydroxyethyl)piperazine, 25 N-methylhomoveratrylamine, nipecotamide, 2-(propylamino)ethanol, 2-(methylamino)ethanol, N-methylhydroxylamine hydrochloride, N,O-dimethylhydroxylamine hydrochloride, Boc-piperazine, 2-aminobenzyl alcohol, 3-aminobenzyl alcohol, 4-aminophenylethyl alcohol, N-benzylaspartic acid, DL-proline, isonipecotic acid, nipecotic acid

30 Step 1:

The series of 24 amines (0.18 mmol) was added to each vial in a 4 x 6 array of tubes. To each was added 0.6 ml of DMF, DIEA (0.091 mL, 0.525 mmol), and chlorotrimethylsilane (0.040 ml, 3.15 mmol), and the array was agitated for 2.5 h.

To a separate solution of (R) methyl (2-pentyl)succinate mono acid (1.0 g, 5.0 35 mmol) in DMF (14 ml) at 0°C was added EDC (0.96 g, 5.0 mmol) and DIEA (209 ml, 1.2

mmol). This mixture was stirred for 20 min, and aliquots (0.42 ml, 0.15 mmol activated acid) were added to each vial in the amine array, followed by a catalytic amount (ca. 5 mg) of DMAP. The mixtures were agitated for 18 h, diluted with 3 ml ethyl acetate, washed with 10% NaHSO₄ (aqueous) and brine, filtered through Na₂SO₄ and evaporated. The
5 resulting methyl esters were placed in a vacuum chamber under high vacuum overnight, weighed, and analyzed by HPLC/MS.

Step 2:

A methanolic solution of NH₂OH HCl (2 M, 15 ml) was cooled to 0°C. A methanolic solution of KOH (3 M, 15 ml) was added and the mixture was stirred for 30 min
10 and filtered. An aliquot (2 ml, 2.0 mmol NH₂OH) of this solution was added to each of the reaction vials containing the methyl ester product of the previous reaction. The array was agitated for 30 min and H⁺ Dowex (220 mg, 1.125 mmol) was added to each vial and the solutions were filtered, evaporated and purified by preparative HPLC. The product-containing fractions were collected, evaporated in a Savant evaporating centrifuge,
15 and re-analyzed by HPLC/MS.

5.6.5 Synthesis of Compounds of Formula XX

Compounds represented by the following general structure are synthesized as described in section 5.6.5. See Scheme III for a general synthetic scheme and Table 16 for
20 representative R groups.

D-Phenylalanine phenyl amide hydrochloride was prepared from Boc-Phe-NHS by standard procedures. This salt (54 mg, 0.20 mmol) was dissolved in 0.8 ml of DCM containing (R) tert-butyl (2-pentyl)succinate mono N-hydroxysuccinimide ester (740 mg, 0.19 mmol) and DIEA (37 ml, 0.21 mmol). The mixture was stirred at 35 °C for 24 h,
25 diluted with DCM and washed once each with 0.2 M HCL, NaHCO₃ (saturated), and brine, filtered through MgSO₄, and evaporated. The residue was analyzed by HPLC/MS and NMR and carried on without further purification.

The succinyl tert-butyl ester (64 mg, 0.137 mmol) was dissolved in 1 ml of 100:5:1 TFA:water:DMS, stirred for 30 min, and evaporated. The residue was then dissolved in 2
30 ml 3:1 benzene:methanol and trimethylsilyldiazomethane in hexanes (2.0 M, 0.50 ml, 1.0 mmol) was added dropwise with stirring. After 20 min the color was discharged by dropwise addition of acetic acid and the solution was evaporated to yield approximately 75 mg of a pale yellow oil which was analyzed by HPLC/MS and carried on without further purification.


35

A methanolic solution of $\text{NH}_2\text{OH HCl}$ (2 M, 0.50 ml) was cooled to 0°C . A methanolic solution of KOH (3 M, 0.5 ml) was added and the mixture was stirred for 30 min and filtered. An aliquot (0.35 ml, 0.35 mmol NH_2OH) of this solution was added to a solution of the succinyl methyl ester from the previous reaction (0.137 mmol theoretical) in 0.2 ml MeOH. The mixture was stirred for 30 min and HCl (12 M, 50 ml, 0.69 mmol) was added followed immediately by 100 ml 1 M triethylammonium bicarbonate (pH 7). The mixture was evaporated and purified by preparative HPLC. The product-containing fractions were collected and evaporated, providing 21 mg (36% yield) of the hydroxamic acid. MS (ES), 393.1, 426.2, 448.1 (M-NHOH, M+H, M+Na); $^1\text{H NMR}$ (CD_3OD) δ 0.93 (t, $J = 7.0$, 3 H), 1.09-1.49 (m, 8 H), 2.23 (dd, $J = 5.1, 15.3$, 1 H), 2.33-2.55 (m, 1 H), 2.65-2.78 (m, 1 H), 2.98 (dd, $J = 10.9, 14.1$, 1 H), 3.50 (dd, $J = 4.4, 14.0$, 1 H), 4.4.70-4.90 (m, 1 H), 7.15-7.60 (m, 10), 7.79 (d, $J = 7.7$, 2 H).

5.7.1 Synthesis of Compounds of Formula XVIII

Compounds represented by the Formula XVIII are synthesized as described in section 5.7.1. See Schemes XVI and XVII for general synthetic schemes and Table 14 for representative R groups.


Synthesis of compound 239. A mixture of 2-amino-4,5,6, 7-tetrahydrobenzo[B] thiophene-3-carboxylic acid ethyl ester (200 mg, 0.89 mmol), 3-nitrobenzenesulfonyl chloride (295 mg, 1.33 mmol), and pyridine (4.4 ml) was heated under reflux for 1 h. The reaction mixture was concentrated in vacuo. CHCl_3 was added to the residue, then washed with 1 N HCl and dried over MgSO_4 . The solvent was removed in vacuo, and the residue was purified by preparative TLC (EtOAc/n-Hexane , 1:4), and recrystallization ($\text{CH}_2\text{Cl}_2/\text{n-Hexane}$) to give compound 239 (66 mg). Analyzed by $^1\text{H NMR}$ (DMSO-d_6) δ : 10.87 (br s, 1H), 8.6-8.5 (m, 2H), 8.23-8.17 (m, 1H), 8.0-7.9 (m, 1H), 4.09 (q, $J=7.0$ Hz, 2H), 2.7-2.5 (m, 4H), 1.8-1.65 (m, 4H), 1.2 (t, $J=7.0$ Hz, 3H). HPLC/MS. Calculated weight 410.47. Found 409.0 (M-H).



Scheme XVI

10

Synthesis of compound 240. Synthesis of compound 240 was completed as described above for compound 239 using 4-nitrobenzenesulfonyl chloride in place of 3-nitrobenzenesulfonyl chloride. See Scheme XVII. Analyzed by ^1H NMR (DMSO- d_6) δ : 10.88 (br s, 1H), 8.51-8.43 (m, 2H), 8.1-8.02 (m, 2H), 4.12 (q, $J=7.0$ Hz, 2H), 2.7-2.55 (m, 4H), 1.85-1.65 (m, 4H), 1.22 (t, $J=7.0$ Hz, 3H). HPLC/MS. Calculated weight 410.47. Found 409.0 (M-H).



Scheme XVII

25 **Synthesis of the NH_2 derivative.** To a solution of compound 240 (250 mg, 0.61 mmol) in EtOH (150 ml) was added 10% Pd/C (150 mg) and it was shook 2 h under H_2 at room temperature. The catalyst was removed by filtration and the filtrate was concentrated in vacuo. Recrystallization from $\text{CHCl}_3/\text{EtOH}/\text{diisopropyl ether}$ gave the NH_2 derivative (83 mg).

30

Synthesis of compound 241. The reaction mixture containing -NH_2 derivative (52 mg, 0.14 mmol), acetic anhydride (56 mg, 0.55 mmol), and CHCl_3 (1 ml) was stirred overnight at room temperature, and concentrated. The residue was purified by chromatography (EtOAc/n-Hexane, 1:2 to 1:1), and recrystallization ($\text{CH}_2\text{Cl}_2/\text{n-Hexane}$) to give compound 241 (28 mg). Analyzed by ^1H NMR (DMSO- d_6) δ : 10.43 (br s, 1H), 10.40 (br s, 1H),

7.84-7.74 (m, 4H), 4.18 (q, J=7.2 Hz, 2H), 2.7-2.55 (m, 4H), 1.82-1.65 (m, 4H), 1.26 (t, J=7.2 Hz, 3H). HPLC/MS. Calculated weight 422.53. Found 377.0, 423.0 (M-EtO, M+H).

- 5 **Synthesis of compound 242.** To a solution of formic acid (19 mg, 0.41 mmol) in CHCl_3 (0.5 ml) was added acetic anhydride (41 mg, 0.40 mmol), followed by the -NH_2 derivative (30 mg, 0.08 mmol). The reaction mixture was stirred 1 h at room temperature, and concentrated. The residue was purified by chromatography (EtOAc/n-Hexane, 1:4 to 1:1), and recrystallization (CH_2Cl_2 / diisopropyl ether) to give compound 242 (7 mg). Analyzed
- 10 by ^1H NMR (DMSO-d_6) δ : 10.69 (br s, 1H), 10.46 (br s, 1H), 8.41 (s, 1H), 7.81 (s, 4H), 4.18 (q, J=7.2 Hz, 2H), 2.75-2.5 (m, 4H), 1.87-1.57 (m, 4H), 1.27 (t, J=7.2 Hz, 3H). HPLC/MS. Calculated weight 408.50. Found 363.0, 409.0, 431.0 (M-EtO, M+H, M+Na).
- 15 **Synthesis of compound 243.** The reaction mixture containing the NH_2 derivative (104 mg, 0.27 mmol), methanesulfonyl chloride (80.3 ml, 1.04 mmol), Et_3N (29.3 ml, 0.21 mmol), and pyridine (1 ml) was heated under reflux for 1 h, and then concentrated. The residue was diluted with CHCl_3 , washed with 1N HCl, dried (MgSO_4) and concentrated. Purification by chromatography (EtOAc/n-Hexane, 1:2) and recrystallization (CH_2Cl_2 / diisopropyl ether)
- 20 gave compound 243 (17 mg). Analyzed by ^1H NMR (DMSO-d_6) δ : 10.7-10.3 (m, 2H), 7.8 (d, J=9.0 Hz, 2H), 7.38 (d, J=9.0 Hz, 2H), 4.18 (q, J=7.1 Hz, 2H), 3.2 (s, 3H), 2.7-2.5 (m, 4H), 1.88-1.6 (m, 4H), 1.27 (t, J=7.1 Hz, 3H). HPLC/MS. Calculated weight 458.58. Found 457.0 (M-H).

25 5.7.4 Synthesis of Compounds of Formula XXI

Compounds represented by the following general structure are synthesized as described in section 5.7.4. See Scheme XVIII for general synthetic schemes and Table 18 for representative R groups.

30 Synthesis of Capped Dipeptide Series


Synthesis of compound 274. Boc-Val-OSu (0.500 g, 1.59 mmol) was added to a solution of H-Nle-OMe.HCl (0.318 g, 1.75 mmol) in DCM (5 ml). DIEA (0.410 g, 0.553 ml, 3.18 mmol) was added and the reaction was shaken overnight. The reaction was diluted with DCM, washed with 5% aqueous citric acid, 5% aqueous K_2CO_3 and brine. The organic

35

layers were dried over Na_2SO_4 and the solvent removed in vacuo. The residue was treated with 4.0 M HCl/dioxane and again the solvent was removed by vacuum.

The resulting amine was dissolved in DCM (2 ml). DIEA (0.172 g, 0.232 ml, 4.02 mmol) was added followed by the addition of benzoyl chloride (0.198 g, 0.164 ml, 1.41 mmol). The reaction was stirred over night. The reaction was diluted with ether, washed with 5% citric acid, 5% K_2CO_3 and brine. The organic layers were dried over Na_2SO_4 , and the solvent removed in vacuo. The residue was purified via flash chromatography to give the methyl ester.

The ester was taken up in a basic solution of hydroxyl amine (1.0 M hydroxylamine; 0.5 M KOH) in MeOH. This was stirred for 15 min and sat. NaHCO_3 was added to quench the reaction. The solution was extracted 3 X DCM. The combined organic layers were washed with brine, dried over Na_2SO_4 and the solvent removed in vacuo. The residue was purified via flash chromatography to give 0.054 g compound 274 which was analyzed by HPLC/MS. Calculated weight 349.43, Found 348.4 (M-H).



Scheme XVIII

Synthesis of compound 275. Synthesis of compound 275 was completed as described above for compound 274 using D-Boc-Val-OSu in place of Boc-Val-OSu. Analyzed by HPLC/MS. Calculated weight 349.43, Found 348.4 (M-H).

Synthesis of compound 276. Synthesis of compound 276 was completed as described above for compound 274 using H-Cha-Ome .HCl in place of H-Nle-OMe.HCl. Analyzed by HPLC/MS. Calculated weight 389.50, Found 388.2 (M-H).

Additional Capped Dipeptides

See scheme XIX.

Step 1:

To a stirring solution of Boc-Nle-OH (0.7 g, 3 mmol) and triethylamine (0.42 ml, 3 mmol) in 1: 1 THF/Ether (15 ml) was added isobutylchloroformate (0.39 ml, 3 mmol) at -20°C. After stirring for 30 min at -20°C, a diazomethane ethereal solution (30 ml, 0.1 M) was added and allowed to stir at -20°C for 30 min and gradually warm to room temperature. After stirring for 16 h, the solution was quenched with a few drop of acetic acid and poured into ether (30 ml), washed with saturated aqueous NaHCO₃ (30 ml) and brine (30 ml). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The crude material was purified by flash chromatography (25% EtOAc/Hexane) to provide 0.25 g (34%) of diazoketone as product. Analyzed by ¹H NMR (CDCl₃) δ: 0.98 (t, 3H), 1.38 (m, 4H), 1.52 (s, 9H), 1.61 (m, 1H), 1.95 (m, 1H), 4.23 (bs, 1H), 5.18 (d, 1H), 5.55 (s, 1H).

Step 2:

To a stirring solution of diazoketone (0.73 g, 3 mmol) in MeOH (5 ml) at 0°C was added silver benzoate (68 mg, 0.3 mmol) and triethylamine (0.84 ml, 6 mmol). The solution was allowed to stir at 0°C for 2 h then concentrated to dryness. The residue was taken up in EtOAc (30 ml) and washed with H₂O (20ml) and 0.5 N HCl (15 ml). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The crude material was purified by flash chromatography (20% EtOAc/Hexane) to provide 0.55 g (72%) of Boc-Nle-methyl ester as product. Analyzed by ¹H NMR (CDCl₃) δ: 0.98 (t, 3H), 1.38 (m, 4H), 1.52 (m, 11H), 2.62 (m, 2H), 3.76 (s, 3H), 3.98 (m, 1H), 4.98 (d, 1H).

Steps 3 and 4:

Boc-Nle-methyl ester (0.55g, 2.1 mmol) was treated with 4N HCl/dioxane (2 ml) for 30 min. The solution was concentrated to dryness, yielding 0.43 g of Nle-methyl ester HCl salt which was used in the next step.


A mixture of EDC (130 mg, 0.67 mmol), HOBt (120 mg, 0.89 mmol) and Bz-Val-OH (150 mg, 0.67 mmol) was stirred in DMF (3 ml) for 15 min. At which time, a solution of b-Nle-methyl ester HCl salt (87 mg, 0.45 mmol) in DMF (2 ml) was added followed by the addition of triethylamine (63 ml, 0.45 mmol). The resulting mixture was stirred for 16 h then poured into saturated aqueous ammonium chloride (15 ml) and EtOAc (10 ml). The aqueous layer was separated and extracted with EtOAc (10 ml). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated. The crude material was purified by flash chromatography (40%

35

EtOAc/Hexane) to provide 130 mg (80%) of the title compound, a mixture of 2 diastereomers. Analyzed by HPLC/MS. Calculated weight 362.47. Found 363.15 (M+H).

Step 5:

A methanolic solution of $\text{NH}_2\text{OH}\cdot\text{HCl}$ (2 M, 5 ml) was cooled to 0°C . A methanolic solution of KOH (3 M, 5 ml) was added dropwise over 10 min, the mixture was stirred for 30 min and filtered. An aliquot (1.5 ml, 1.5 mmol) of this solution was added to the methyl ester (60 mg, 0.16 mmol). The mixture was allowed to stir for 20 min and acidified to $\text{pH} = 5$ by Dowex acidic resin. The solvent was removed and the residue was purified by preparative HPLC to isolated compound 277 and compound 278 (diastereomers) in pure form. Analyzed by HPLC/MS. Calculated weight 363.46. Found 362.15 (M-H).



Scheme XIX

Step 6:

Methyl ester (60 mg, 0.16 mmol) from step 4 was treated with 1M LiOH/MeOH (2 ml) for 18 h. Dowex acidic resin was added to acidify the solution to pH = 5. The filtrate was collected and concentrated to dryness to give 45 mg (79%) of compound 279 as desired product. Analyzed by HPLC/MS. Calculated weight 348.45. Found 347.10 (M-H).

5.7.2 Synthesis of Compounds of Formula XVII**Synthesis of Capped Valine series**

Compounds represented by Formulas XVI and XVII are synthesized as described in section 5.7.2 See scheme XX for a general synthetic scheme and Tables 12 and 13 for representative structures.


Synthesis of compound 224. Pentylamine (1.31 g, 1.74 ml, 15.0 mmol) was added to a suspension of Argopore MB-CHO resin (1.0 g, 0.75 mmol) in DCM (5 ml) and trimethylorthoformate (5 ml) and shaken for 3 hr. The resin was filtered and washed quickly 3 times with dry DCM. The resin was then suspended in DCM (5 ml). Sodium triacetoxyborohydride (3.18 g, 15 mmol) was then added and the reaction was shaken over night. The resin was then filtered and washed repeatedly (3 X DMF, 1 X DCM, 1 X MeOH, 3 X DCM) and dried by vacuum to give the pentyl amine resin.

A mixture of Fmoc-Val-OH (0.764 g, 2.25 mmol), PyBOP (1.17 g, 2.25 mmol) and HOBT (0.304 g, 2.25 mmol) in DCM (5 ml) was shaken for 30 min and added to the pentylamine resin (1 g, 0.75 mmol). This was shaken over night and filtered. The resin was washed (3 X DMF, 1 X DCM, 1 X MeOH, 3 X DCM) and dried in vacuo to give the valine capped pentyl amine resin. This resin was treated for 30 min with 20% piperidine in toluene for 30 minutes, washed repeatedly (3 X DMF, 1 X DCM, 1 X MeOH, 3 X DCM) and dried in vacuo to give the deprotected amino acid bound resin.

Diisopropyl carbodiimide (0.473 g, 0.550 ml, 3.75 mmol) was added to 3-bromomethylbenzoic acid (0.806 mg, 3.75 mmol) and DMAP (0.0046 g, 0.0375 mmol) in DMF (3 ml). This solution was shaken for 5 min and added to the above resin (0.500 g, 0.375 mmol) and shaken over night. The resin was washed repeatedly (3 X DMF, 1 X DCM, 1 X MeOH, 3 X DCM) and dried in by vacuum to give the capped valine resin.

1,2,3,4-Tetrahydro-1-naphthylamine (0.057 ml, 0.375 mmol) was added to a suspension of the capped valine resin (0.050 g, 0.0375 mmol) in DMF (1 ml) and shaken overnight. The resin was then filtered and washed repeatedly (3 X DMF, 1 X DCM, 1 X

MeOH, 3 X DCM) to give the compound 224 bound resin. The damp resin was treated with 80% trifluoroacetic acid in DCM for 1 h and filtered. The resin was washed with DCM and the combined filtrate was dried in vacuo to give 4.4 mg of compound 224 which was analyzed by HPLC/MS. Calculated weight 449.64, Found 450 (M+H).



Synthesis of compound 225. Synthesis of compound 225 was completed as described above for compound 224 using butylamine in place of 1,2,3,4-tetrahydro-1-naphthylamine. Analyzed by HPLC/MS. Calculated weight 375.56, Found 376.30 (M+H).

Synthesis of compound 226. Synthesis of compound 226 was completed as described above for compound 224 using phenethylamine in place of 1,2,3,4-tetrahydro-1-naphthylamine. Analyzed by HPLC/MS. Calculated weight 423.60, Found 424.3 (M+H).

Synthesis of compound 227. Synthesis of compound 227 was completed as described above for compound 224 using cyclopentylamine in place of 1,2,3,4-tetrahydro-1-naphthylamine. Analyzed by HPLC/MS. Calculated weight 387.57, Found 388.3 (M+H).

5

Synthesis of compound 228. Synthesis of compound 228 was completed as described above for compound 224 using tetrahydrofurfurylamine in place of 1,2,3,4-tetrahydro-1-naphthylamine. Analyzed by HPLC/MS. Calculated weight 403.57, Found 404.3 (M+H).

10

Synthesis of compound 229. Synthesis of compound 229 was completed as described above for compound 224 using 4-chlorobenzylamine in place of 1,2,3,4-tetrahydro-1-naphthylamine. Analyzed by HPLC/MS. Calculated weight 444.02, Found 444.2 (M+H).

15

Synthesis of compound 230. Synthesis of compound 230 was completed as described above for compound 224 using 4-bromomethylbenzoic acid in place of 3-bromomethylbenzoic acid. Analyzed by HPLC/MS. Calculated weight 449.64, Found 450 (M+H).

20

Synthesis of compound 231. Synthesis of compound 231 was completed as described above for compound 225 using 4-bromomethylbenzoic acid in place of 3-bromomethylbenzoic acid. Analyzed by HPLC/MS. Calculated weight 375.56, Found 376.30 (M+H).

25

Synthesis of compound 232. Synthesis of compound 232 was completed as described above for compound 226 using 4-bromomethylbenzoic acid in place of 3-bromomethylbenzoic acid. Analyzed by HPLC/MS. Calculated weight 423.60, Found 424.3 (M+H).

30

Synthesis of compound 233. Synthesis of compound 233 was completed as described above for compound 227 using 4-bromomethylbenzoic acid in place of 3-bromomethylbenzoic acid. Analyzed by HPLC/MS. Calculated weight 387.57, Found 388.3 (M+H).


35

Synthesis of compound 234. Synthesis of compound 234 was completed as described above for compound 228 using 4-bromomethylbenzoic acid in place of 3-bromomethylbenzoic acid. Analyzed by HPLC/MS. Calculated weight 403.57, Found 404.3 (M+H).


Synthesis of compound 235. Synthesis of compound 235 was completed as described above for compound 229 using 4-bromomethylbenzoic acid in place of 3-bromomethylbenzoic acid. Analyzed by HPLC/MS. Calculated weight 444.02, Found 444.2 (M+H).

5.6.1 Synthesis of Compounds of Formula XXI

Compounds represented by the following general structure are synthesized as described in section 5.6.1. See Scheme XXI for general synthetic scheme and Table 19 for representative R groups.



Scheme XX



Scheme XXII

Synthesis of compound 205

15

Synthesis of triethyl 5-methyl-1,2,2-hexanetricarboxylate. Sodium metal (467 mg, 20.3 mmol) was added to a stirred solution of triethyl 1,1,2-ethane tricarboxylate (5.0 g, 4.7 ml, 20.3 mmol) in ethanol (35 ml). After the metal had dissolved 1-bromo-3-methyl butane (3.1 g, 2.4 ml, 20.3 mmol) was added and the mixture was refluxed for 24 h. The mixture
 20 was cooled, filtered and resulting filtrate was concentrated. The residue was taken up in ether and washed with water. The water layer was back extracted with ether and the combine organic layers were dried (MgSO_4), concentrated and purified by flash chromatography using 10% EtOAc/hexanes to give 5.6 g (87%) of triethyl 5-methyl-1,2,2-hexanetricarboxylate as a colorless oil. The material was analyzed by ^1H
 25 NMR.

Synthesis of 3-carboxy-6-methyl heptanoic acid. Triethyl 5-methyl-1,2,2-hexanetricarboxylate (5.6 g, 17.7 mmol) was added to 55 ml conc. HCl and heated to reflux for 48 h. A white solid precipitate was noted. The reaction mixture was poured into
 30 ice and extracted with ether. The combined extracts were dried (MgSO_4) and concentrated to form a yellow oil. The oil crystallized upon standing to give 2.5 g (92%) of 3-carboxy-6-methyl heptanoic acid as a colorless crystalline solid which was analyzed by ^1H NMR.

35 **Synthesis of isopentyl succinic anhydride.** 3-Carboxy-6-methyl heptanoic acid (2.8 g,

14.9 mmol) was added to 5.2 ml acetyl chloride and heated to reflux for 3 h. Volatile material was removed in vacuo to give 2.5 g crude isopentyl succinic anhydride which was analyzed by ¹H NMR and used without further purification.

- 5 **Synthesis of N-Boc-Valine homopiperidine amide.** Homopiperidine (3.3 g, 33.2 mmol) was added to a solution of N-Boc-Valine (6.0 g, 27.7 mmol) and EDC (6.3 g, 33.2 mmol) in DCM (60 ml). The mixture was stirred for 16 h. The reaction mixture was diluted with DCM, washed with water, brine, dried (MgSO₄) and concentrated. Residue was purified by flash chromatography using 60% EtOAc in hexanes to give 3.4 g N-Boc-Valine
10 homopiperidine amide as a colorless oil. This material analyzed by ¹H NMR.

- Coupling of isopentyl succinic anhydride and isopentyl succinic anhydride to form 1.** TFA (3 ml) was added to a solution of N-Boc-Valine (0.650 g, 2.2 mmol) homopiperidine amide in DCM (12 ml). The ice bath was removed and solution stirred for 4.5 h. Solvent
15 was removed in vacuo and replaced with DCM (20 ml). 0.341 g (2.6 mmol) DIEA was added to the solution followed by addition of isopentyl succinic anhydride (0.442 g, 2.6 mmol). The solution was stirred for 72 h. The mixture was diluted with DCM, washed with brine, dried (MgSO₄) and concentrated. The residue was purified by flash chromatography using EtOAc. 0.740 g (91%) of product 1 was obtained as a colorless oil
20 which was analyzed by ¹H NMR.

- Synthesis of compound 205.** Coupling product 1 (0.740 g, 2.0 mmol) from the previous reaction was added to 0.71 ml acetyl chloride and heated to reflux for 3 h. EtOAc was added and washed with saturated NaHCO₃ (x2), brine, dried (MgSO₄) and concentrated. A
25 cold solution of 0.264 g (3.8 mmol) hydroxamic acid hydrochloride salt in MeOH was treated with a solution of 0.319 g (5.7 mmol) KOH and let stand for 15 min at 0°C. The resulting solution was filtered, added to the residue from the previous step and stirred at room temperature for 48 h. The pH of the solution was neutralized to pH ~ 6 by addition of Dowex 50W-X8 resin. The solution was filtered and concentrated. The residue was
30 purified by flash chromatography using 80% EtOAc in hexanes. Material was analyzed by ¹H NMR and HPLC/MS. Calculated weight 383.25, Found 349.2, 382.3, 406.2 (M-NHOH, M+H, M+Na).

- Synthesis of compound 206.** Synthesis of compound 206 was completed as described
35 above using bromomethyl cyclohexane in place of 1-bromo-3-methyl butane.

Synthesis of compound 207. Synthesis of compound 207 was completed as described above using 1-bromo-trans-2-butene in place of 1-bromo-3-methyl butane.

Synthesis of compound 208. Synthesis of compound 208 was completed as described
5 above using 1-bromobutane in place of 1-bromo-3-methyl butane.

Synthesis of compound 209. Synthesis of compound 209 was completed as described above using benzyl bromide in place of 1-bromo-3-methyl butane.

10 **Synthesis of compound 274.** Boc-Val-OSu (0.5000 g, 1.59 mmol) was added to a solution of H-Nle-OMe.HCL (0.318 g, 1.75 mmol) in DCM (5 ml). DIEA (0.410 g, 0.553 ml, 3.18 mmol) was added and the reaction was agitated overnight. The reaction was diluted with DCM, washed with 5% citric acid, 5% K₂CO₃ and brine. The organic layers were dried over Na₂SO₄ and the solvent removed in vacuo. The residue was treated with 4.0 M
15 HCl/dioxane and again the solvent was removed in vacuo.

The resulting amine was dissolved in DCM (2 ml). DIEA (0.172 g, 0.232 ml, 4.02 mmol) was added followed by the addition of benzoyl chloride (0.198 g, 0.164 ml, 1.41 mmol). The reaction was stirred overnight. The reaction was diluted with ether, washed with 5% citric acid, 5% K₂CO₃ and brine. The organic layers were dried over Na₂SO₄, and
20 the solvent removed in vacuo. The residue was purified via flash chromatography to give the methyl ester. The ester was taken up in a basic solution of hydroxyl amine (1.0 M hydroxylamine; 0.5 M KOH) in MeOH. This was stirred for 15min and saturated NaHCO₃ was added to quench the reaction. The solution was extracted 3 X DCM. The combined organic layers were washed with brine, dried over Na₂SO₄ and the solvent removed in
25 vacuo. The residue was purified via flash chromatography to give 0.054 g compound 274 which was analyzed by HPLC/MS. Calculated weight 349.43, Found 348.4 (M-H).

Synthesis of compound 275. Synthesis of compound 275 was completed as described above for compound 274 using D-Boc-Val-OSu in place of Boc-Val-OSu. Analyzed by
30 HPLC/MS. Calculated weight 349.43, Found 348.4 (M-H).

Synthesis of compound 276. Synthesis of compound 276 was completed as described above for compound 274 using H-Cha-Ome.HCl in place of H-Nle-OMe.HCl. Analyzed by HPLC/MS. Calculated weight 389.50, Found 388.2 (M-H).

35

5.8.1 Synthesis of Hydantoin Derivatives

Compounds represented by Formulas XIX and XX are synthesized as described in section 5.8.1. See scheme XXIII for a general synthetic scheme and Tables 15-16 for representative structures.

5 **Synthesis of compound 262.**

Step 1:

D,L-Leucine (6.6 g, 0.050 mol) and KOCN (4.9 g, 0.060 mol) were allowed to react in 30 ml of water and 25 ml of pyridine for 30 min at 60°C. The solution mixture was cooled to room temperature then extracted with EtOAc (2 X 150 ml). 6N HCl (30 ml) and
10 acetic acid (30 ml) was added to the aqueous phase and the mixture was heated to reflux for another 30 min. Upon cooling to room temperature, white solid precipitate out from solution. The solid was collected by vacuum filtration and washed repeatedly with water and ether. Yield 6.56 g, 84% of desired product. Analyzed by NMR. ¹H NMR (300 MHz, DMSO- d₆) δ: 0.89 (dd, 6H), 1.38-1.49 (m, 2H), 1.76 (m, 1H), 4.00 (dd, 1H), 8.02 (s, 1H),
15 10.60 (s, 1H).

Step 2:

A mixture of isobutylhydantoin (468 mg, 3 mmol), benzyl bromide (392 ml, 3.3 mmol) and K₂CO₃ (622 mg, 4.5 mmol) in 7 ml of acetone were heated at 50°C overnight. EtOAc (15 ml) was added to the resulting mixture and washed with water then brine. The
20 organic phase was dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The crude product was triturated with 40% DCM/hexane to give 420 mg of white solid as product. Yield 57%. Analyzed by NMR. ¹H NMR (300 MHz, CDCl₃) δ: 0.99 (dd, 6H), 1.53 (m, 1H), 1.65 (m, 2H), 4.00 (dd, 1H), 4.68 (s, 2H), 6.02 (bs, 1H), 7.29-7.42 (m, 5H).


25 Step 3:

A mixture of isobutyl-benzyl-hydantoin (246 mg, 1 mmol), methylbromoacetate (114 ml, 1.2 mmol) and K₂CO₃ (207 mg 1.5 mmol) in 3 ml of acetone were heated in a seal tube at 50°C overnight. EtOAc (10 ml) was added to the resulting mixture and washed with water then brine. The organic phase was dried over anhydrous sodium sulfate and
30 concentrated to dryness. The crude oil was carried to next step without further purification.

Step 4:

A methanolic solution of NH₂OH.HCl (2.78 g in 34 ml) was cooled to 0°C. A methanolic solution of NaOH (3 g in 6 ml) was added, the mixture was stirred for 30 min and filtered. An aliquot (1 ml, 1.0 mmol NH₂OH) of this solution was added to a vial
35 containing hydantoin methyl ester product (83 mg, 0.26 mmol) of the previous reaction.

The vial was shaken for 15 min then acidified to pH=5-6 with Dowex H⁺ resin.. The solution was then filtered and concentrated to dryness. The crude material was purified by flash chromatography using 5% MeOH/DCM. Analyzed by HPLC/MS. Calculated molecular weight 319.36. Found 320.1 (M+H).



Synthesis of compound 261. Synthesis of compound 261 was completed as described above for compound 262 using D,L-Val in place of D,L-Leu. Calculated molecular weight 305.33. Found 306.1 (M+H).

Synthesis of compound 263. Synthesis of compound 263 was completed as described above for compound 262 using D,L-Phe in place of D,L-Leu. Calculated molecular weight 353.38. Found 354.1 (M+H).

Synthesis of compound 256. Synthesis of compound 256 was completed as described above for compound 262 using D,L-Nle in place of D,L-Leu. Analyzed by NMR. ¹H NMR (300 MHz, CD₃OD) δ: 0.88 (t, 3H), 1.06 (m, 1H), 1.29 (m, 3H), 1.88 (m, 2H), 3.79 (m, 1H), 4.25 (m, 2H), 4.70 (m, 2H), 7.35 (m, 5H).

Synthesis of compound 255. Synthesis of compound 255 was completed as described above for compound 262 using D,L-Val in place of D,L-Leu and iodomethane in place of benzyl bromide. Analyzed by NMR. ¹H NMR (300 MHz, CD₃OD) δ: 0.98 (dd, 6H), 2.27 (m, 1H), 3.00 (s, 3H), 4.04 (d, 1H), 4.03 (dd, 2H).

Synthesis of compound 258. Synthesis of compound 258 was completed as described

above for compound 262 using iodomethane in place of benzyl bromide. Analyzed by NMR. ¹H NMR (300 MHz, CD₃OD) δ: 0.97 (dd, 6H), 1.73 (m, 2H), 1.90 (m, 1H), 3.00 (s, 3H), 4.05 (dd, 2H), 4.16 (dd, 1H).

- 5 **Synthesis of compound 260.** Synthesis of compound 260 was completed as described above for compound 262 using D,L-Nle in place of D,L-Leu and iodomethane in place of benzyl bromide. Calculated molecular weight 243.26. Found 244 (M+H).

- Synthesis of compound 257.** Synthesis of compound 257 was completed as described
10 above for compound 262 using D,L-Met in place of D,L-Leu and iodomethane in place of benzyl bromide. Analyzed by HPLC/MS. Calculated molecular weight 261.30. Found 262 (M+H).

- Synthesis of compound 259.** Synthesis of compound 259 was completed as described
15 above for compound 262 using D,L-Phe in place of D,L-Leu and iodomethane in place of benzyl bromide. Analyzed by HPLC/MS. Calculated molecular weight 277.28. Found 278.1 (M+H).

Synthesis of compound 300 (Scheme XXIV)

20 Step 1:

- Chlorotrimethylsilane (274 ml, 2.17 mmol) was added to a stirring solution of D,L-Phenylalanine (358 mg, 2.17 mmol) and triethylamine (302 ml, 2.17 mmol) in anhydrous DCM (6 ml). The resulting mixture was allowed to stir for 15 min then ethyl 2-isocyanato-4-(methylthio)butyrate (400 mg, 1.97 mmol) was added. After stirring for 18
25 h, 100 ml of TFA was added and allowed to stir for additional 30 min then evaporated to dryness. The residue was taken in EtOAc (6 ml) and extracted with 1N HCl (2 X 3 ml) then brine. The organic phase was dried over anhydrous sodium sulfate, filtered and solvent removed under reduced pressure. The crude product was carried on to next step without further purification.

30 Step 2:


- A mixture of acid (830 mg, 2.25 mmol) obtained from step 1 and 1,1'-carbonyldiimidazole (504 mg, 3.11 mmol) in 10 ml of anhydrous THF was allowed to stir at room temperature overnight. EtOAc (15 ml) was added to the mixture and washed with 1N HCl (10 ml), saturated aqueous NaHCO₃ (10 ml), and brine. The organic phase
35 was dried over anhydrous sodium sulfate and concentrated to dryness. The crude product

was purified by flash chromatography using 25% EtOAc/Hexane as eluting solvent.

Step 3:

A methanolic solution of $\text{NH}_2\text{OH}\cdot\text{HCl}$ (2.78 g in 34 ml) was cooled to 0°C . A methanolic solution of NaOH (3 g in 6 ml) was added, the mixture was stirred for 30 min and filtered.

- 5 An aliquot (1 ml, 1.0 mmol NH_2OH) of this solution was added to a vial containing hydantoin methyl ester product (57 mg, 0.16 mmol) of the previous reaction. The vial was shaken for 30 min then acidified to pH 5-6 with Dowex H^+ resin. The solution was then filtered and concentrated to dryness. The crude material was purified by flash chromatography using 5% MeOH/DCM. Analyzed by HPLC/MS. Calculated molecular weight 337.40. Found 336.1 (M-H).



Scheme XXIV

Synthesis of compound 246. Synthesis of compound 246 was completed as described above for compound 300 using D,L-Val in place of D,L-Phe. Analyzed by HPLC/MS.

- 30 Calculated molecular weight 289.36. Found 288.1 (M-H).

Synthesis of compound 244. Synthesis of compound 244 was completed as described above for compound 300 using D,L-Leu in place of D,L-Phe. Analyzed by HPLC/MS. Calculated molecular weight 303.38. Found 302.1 (M-H).

35

Synthesis of compound 248. Synthesis of compound 248 was completed as described above for compound 300 using D,L-Nle in place of D,L-Phe. Analyzed by HPLC/MS. Calculated molecular weight 303.38. Found 302.1 (M-H).

- 5 **Synthesis of compound 245.** Synthesis of compound 245 was completed as described above for compound 300 using D,L-Met in place of D,L-Phe. Analyzed by HPLC/MS. Calculated molecular weight 321.42. Found 320.05 (M-H).

- 10 **Synthesis of compound 253.** Synthesis of compound 253 was completed as described above for compound 300 using ethyl 2-isocyanato-4-methylvalerate in place of ethyl 2-isocyanato-4-(methylthio)butyrate. Analyzed by HPLC/MS. Calculated molecular weight 321.42. Found 320.05 (M-H).

- 15 **Synthesis of compound 247.** Synthesis of compound 247 was completed as described above for compound 300 using ethyl 2-isocyanato-3-phenylpropionate in place of ethyl 2-isocyanato-4-(methylthio)butyrate. Analyzed by HPLC/MS. Calculated molecular weight 321.42. Found 320.05 (M-H).

- 20 **Synthesis of compound 252.** Synthesis of compound 252 was completed as described above for compound 300 using D,L-Met in place of D,L-Phe and ethyl 2-isocyanato-4-methylvalerate in place of ethyl 2-isocyanato-4-(methylthio)butyrate. Analyzed by HPLC/MS. Calculated molecular weight 303.38. Found 302.1 (M-H).

- 25 **Synthesis of compound 254.** Synthesis of compound 254 was completed as described above for compound 300 using D,L-Nle in place of D,L-Phe and ethyl 2-isocyanato-4-methylvalerate in place of ethyl 2-isocyanato-4-(methylthio)butyrate. Analyzed by HPLC/MS. Calculated molecular weight 285.34. Found 284.1 (M-H).

- 30 **Synthesis of compound 251.** Synthesis of compound 251 was completed as described above for compound 300 using D,L-Val in place of D,L-Phe and ethyl 2-isocyanato-3-methylbutyrate in place of ethyl 2-isocyanato-4-(methylthio)butyrate. Analyzed by HPLC/MS. Calculated molecular weight 257.29. Found 256.1 (M-H).

Synthesis of compound 301 (Scheme XXV)Step 1:

EDC (1.03 g, 5.39 mmol) was added to a solution of Boc-Val (1.17 g, 5.39 mmol) in 8 ml of DCM and stirred for 30 min. At which time, pyrrolidine (0.45 ml, 7.42 mmol) was added and allowed the resulting mixture to stir for 18 h. After evaporation of solvent, the residue was taken up in EtOAc (10 ml) and extracted with 1N HCl (5 ml), saturated aqueous NaHCO₃ (5 ml) then brine (5 ml). The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to give crude Boc amide.

Steps 2 and 3:

Boc amide (0.9 g, 3.3 mmol) was treated with 9 ml of 20% TFA in DCM for 3 h. The volatile material W-S removed b vacuum to give the TFA salt of the amino amide. To a solution of amino amide TFA salt (0.84 g, 2.9 mmol) and triethylamine (835 ml, 5.8 mmol) in 4 ml anhydrous DCM was added 2-isocyanato-4-methylvalerate (447 ml, 2.4 mmol) and stirred at room temperature overnight. After removal of volatile material, the residue was taken up in EtOAc (10 ml) and washed with 0.5N HCl (5 ml), water then brine. The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The crude was purified by flash chromatography using 75% EtOAc/hexane as eluting solvent. Analyzed by HPLC/MS. Calculated weight 355.48. Found 356.20 (M+H).

Step 4:

0.5 N NaOMe (1 ml, in MeOH) was added to a vial containing urea (0.17 g, 0.5 mmol) obtained from previous step. The mixture was allowed to shake for 6 h then acidified to pH 5-6 with Dowex H⁺ resin. The solution was then filtered and concentrated to dryness. The crude material was purified by flash chromatography using 75% EtOAc/hexane. Analyzed by NMR. ¹H NMR (300 MHz, CDCl₃) δ: 0.96-1.10 (m, 12H), 1.59 (m, 1H), 1.82-2.06 (m, 6H), 3.15 (m, 1H), 3.56 (t, 4H), 4.12 (d, 1H), 4.39 (d, 1H), 6.70 (d, 1H).


Step 5:

A mixture of hydantoin (126 mg, 0.41 mmol), methylbromoacetate (154 ml, 1.6 mmol) and K₂CO₃ (113 mg 0.82 mmol) in 4 ml of acetone were heated at 60°C for 24 h. After evaporation of solvent, EtOAc (15 ml) was added to the residue and washed with water then brine. The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The crude material was purified by flash chromatography using 50% EtOAc/hexane.

Step 6:

A methanolic solution of $\text{NH}_2\text{OH}\cdot\text{HCl}$ (2.78 g in 34 ml) was cooled to 0°C . A methanolic solution of NaOH (3 g in 6 ml) was added, the mixture was stirred for 30 min and filtered. An aliquot (0.4 ml, 0.4 mmol NH_2OH) of this solution was added to a vial containing hydantoin methyl ester product (24 mg, 0.06 mmol) of the previous reaction.

- 5 The vial was shaken for 15 min then acidified to pH 5-6 with Dowex H^+ resin.. The solution was then filtered and concentrated to dryness. The crude material was purified by flash chromatography using 5% MeOH/DCM . Analyzed by HPLC/MS. Calculated molecular weight 382.46. Found 381.2 (M-H).



- Synthesis of compound 266.** Synthesis of compound 266 was completed as described above for compound 301 using Boc-Gly in place of Boc-Val. Analyzed by HPLC/MS. Calculated molecular weight 340.38. Found 339.15 (M-H).

- Synthesis of compound 267.** Synthesis of compound 267 was completed as described above for compound 301 using Boc-Gly in place of Boc-Val and hexamethyleneimine in place of pyrrolidine. Analyzed by HPLC/MS. Calculated molecular weight 368.43. Found 367.2 (M-H).

- Synthesis of compound 268.** Synthesis of compound 268 was completed as described above for compound 301 using hexamethyleneimine in place of pyrrolidine. Analyzed by HPLC/MS. Calculated molecular weight 410.51. Found 409.20 (M-H).

Synthesis of compound 269. Synthesis of compound 269 was completed as described above for compound 301 using 2-isocyanato-3-propylbutyrate in place of 2-isocyanato-4-methylvalerate. Analyzed by HPLC/MS. Calculated molecular weight 410.51. Found 409.20 (M-H).

5

Synthesis of compound 272 (Scheme XXVI)

Step 1:

A mixture of Boc-Phe-Osu (1.09 g, 2.8 mmol) and R-(-)-Prolinol (0.5 g, 4.9 mmol) was stirred in 5 ml THF for 18 h.. After evaporation of solvent, the residue was taken up in EtOAc (10 ml) and extracted with 1N HCl (5 ml), saturated aqueous NaHCO₃ (5 ml) then brine (5 ml). The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to give crude Boc amide.

10

Steps 2 and 3:

Same as step 2 and step 3 in the synthesis of compound 301.

15

Step 4:

Same as step 4 in the synthesis of compound 301.

Step 5:

1M NaHMDS (139 ml, 0.14 mmol) was added to a solution of hydantoin (47.3 mg, 0.14 mmol) in 1 ml of THF at -78°C. The solution was allowed to warm to 0°C then added chlorotrimethylsilane (17.6 ml, 0.14 mmol). After stirring for 15 min, another equivalent of NaHMDS (139 ml, 0.14 mmol) was added at 0°C followed by the addition of methylbromoacetate (19.7 ml, 0.21 mmol). The resulting mixture was allowed to warm to room temperature and stir for overnight. After evaporation of solvent, EtOAc (10 ml) was added to the residue and washed with water then brine. The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The crude material was purified by flash chromatography using 75% EtOAc/hexane.

20

25


Step 6:

Same as step 6 in the synthesis of compound 301.

Analyzed by HPLC/MS. Calculated molecular weight 460.53. Found 461.2 (M+H).

30

35



Scheme XXVI

Synthesis of compound 273. Synthesis of compound 273 was completed as described above for compound 272 using Boc-D-Phe-OSu in place of Boc-Phe-OSu. Analyzed by HPLC/MS. Calculated molecular weight 460.53. Found 461.2 (M+H).

25

Synthesis of compound 270. Synthesis of compound 270 was completed as described above for compound 272 using Boc-Gly-OSu in place of Boc-Phe-OSu. Analyzed by HPLC/MS. Calculated molecular weight 370.41. Found 371 (M+H).

30

Synthesis of compound 271. Synthesis of compound 271 was completed as described above for compound 272 using Boc-Val-OSu in place of Boc-Phe-OSu. Analyzed by HPLC/MS. Calculated molecular weight 412.49. Found 413 (M+H).

35

Synthesis of compound 264 (Scheme XXVII)Step 1:

t-Butyl bromoacetate (1.34 ml, 9.1 mmol) was added to a suspension solution of L-Leucine-methylester (3.3 g, 18.1 mmol) and triethylamine (2.52 ml, 18.1 mmol) in 40 ml DMF and stirred for 18 h. The resulting mixture was poured into 40 ml of saturated aqueous ammonium chloride and extracted with ether (2 X 50 ml). The combined ethereal layers were dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The crude material was purified by flash chromatography using 20% EtOAc/hexane. Yield 2.3 g, 98%. Analyzed by NMR. ¹H NMR (300 MHz, CDCl₃) δ: 1.00 (d, 6H), 1.53 (s, 9H), 1.58 (m, 2H), 1.82 (m, 1H), 3.32-3.40 (m, 3H), 3.78 (s, 3H).

Step 2:

A mixture of amine (1.6 g, 6.20 mmol) and methyl-3-isocyanato benzoate (1 g, 5.64 mmol) was stirred in 10 ml of DCM for 18 h. After removal of volatile material, the residue was taken up in EtOAc (30 ml) and washed with 1 N HCl (15 ml) then brine. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The crude material was carried to next step without purification.

Step 3:

Triethylamine (0.8 ml, 5.7 mmol) was added to a solution of urea (2.4 g, 5.64 mmol) obtained from previous step and heated to reflux in 10 ml of THF for 14 h. Volatile material was then removed in vacuo and the residue was purified by flash chromatography using 30% EtOAc/Hexane. Analyzed by NMR. ¹H NMR (300 MHz, CDCl₃) δ: 1.05 (dd, 6H), 1.57 (s, 9H), 1.82-2.03 (m, 3H), 3.80 (d, 1H), 4.00 (s, 3H), 4.44 (dd, 1H), 4.65 (d, 1H), 7.64 (t, 1H), 7.73 (d, 1H), 8.11 (d, 1H), 8.17 (s, 1H).


Step 4:

Hydantoin (0.2 g, 0.5 mmol) was treated with 50% TFA/DCM (3 ml) for 16 h. The volatile material was removed in vacuo to yield acid as crude product. Analyzed by NMR. ¹H NMR (300 MHz, CDCl₃) δ: 1.05 (dd, 6H), 1.92-2.03 (m, 3H), 3.94 (d, 1H), 4.00 (s, 3H), 4.44 (dd, 1H), 4.73 (d, 1H), 7.64 (t, 1H), 7.73 (d, 1H), 8.15 (d, 1H), 8.19 (s, 1H).

Step 5:

EDC (100 mg, 1.04 mmol) and HOBt (70 mg, 1.04 mmol) was added to a solution of acid (180 mg, 0.52 mmol) in DMF (3 ml) and stirred for 1 h. O-Trimethylsilylhydroxylamine (90 ml, 1.04 mmol) was then added and the solution was stirred for 16 h. Saturated ammonium chloride solution was added and the mixture was extracted with EtOAc. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The crude material was purified by flash chromatography using 5% MeOH/DCM.

Analyzed by HPLC/MS. Calculated molecular weight 363.37. Found 362.10 (M-H).



Scheme XXVII

Synthesis of compound 265. Synthesis of compound 265 was completed as described above for compound 264 using 3-acetylphenylisocyanate in place of methyl-3-isocyanato benzoate. Analyzed by HPLC/MS. Calculated molecular weight 362.39. Found 4361.10 (M-H).

Synthesis of Compounds 310-311 (Scheme XXVIII)

Step 1:

Lithium diisopropylamide (3 ml, 6 mmol, 2M in hexane) was added to a solution of S-(+)-trityloxymethyl-butylolactone (1.79 g, 5 mmol) in 15 ml THF at -78°C . After stirring for 30 min at -78°C , a solution of crotyl bromide (0.66 ml, 5.5 mmol) in 5 ml THF was added. The solution mixture was allowed to stir for 4 h at -78°C and then quenched with aqueous ammonium chloride (10 ml). After warming to room temperature, the resulting solution was extracted with EtOAc (2 X 30 ml). The combined organic layers were dried over anhydrous sodium sulfate and concentrated to dryness. The residue was purified by flash chromatography (25% EtOAc/Hexane) to give desired product (1.7g, 84%). Analyzed by ^1H NMR (CDCl_3) δ : 1.38 (m, 1H), 1.7 (m, 3H), 2.05-2.18 (m, 2H), 2.60 (m, 1H), 2.94 (m, 1H), 3.21 (dd, 1H), 3.48 (dd, 1H), 4.66 (m, 1H), 5.43 (m, 1H), 5.62 (m, 1H), 7.24-7.63 (m, 15H).

Step 2:

A mixture of (S)- -buten-2-yl- -trityloxymethyl- -butyrolactone (1.7 g, 4.13 mmol), 10% Pd/C (228 mg) and EtOAc (80 ml) was hydrogenated at 45 psi for 3 h. The Pd catalyst was removed by filtration through a celite pad and washed with EtOAc (40 ml). The filtrate was concentrated to give 1.68 g (99%) of title compound. Analyzed by HPLC/MS.

5 Calculated weight 414. Found 437 (M+Na).

Step 3:

Lithium diisopropylamide (2.43 ml, 4.86 mmol, 2 M in hexane) was added to a solution of (S)- -butyl- -trityloxymethyl- -butyrolactone (1.69 g, 4.05 mmol) in 20 ml THF at -78°C. After stirring for 1 h at -78°C, a solution of allyl bromide (0.45 ml, 5.26 mmol) in 2 ml THF was added. The solution mixture was allowed to stir for 4 h at -78°C and then quenched with aqueous ammonium chloride (10 ml). After warming to room temperature, the resulting solution was extracted with EtOAc (2 X 30 ml). The combined organic layers were dried over anhydrous sodium sulfate and concentrated to dryness. The crude residue was carried to next step without purification. Yield 1.8 g, quantitative. Analyzed by ¹H NMR (CDCl₃) δ: 0.98 (t, 3H), 1.38 (m, 4H), 1.75(m, 2H), 2.05 (m, 2H), 2.42 (m, 2H), 3.25 (dd, 1H), 3.42 (dd, 1H), 4.60 (m, 1H), 5.23 (dd, 2H), 5.95 (m, 1H), 7.24-7.63 (m, 15H).

Step 4:

Trifluoroacetic acid (10 ml) was added to a solution of [S-(R*,R*)]- -butyl- -(propen-2-yl)- -trityloxymethyl- -butyrolactone (1.8 g, 3.96 mmol) in DCM (30 ml) at 0°C. The solution was stirred for 35 min at 0°C then concentrated to dryness. Saturated aqueous sodium chloride solution (30 ml) was poured into residue and extracted with DCM (2 x 20 ml). The combined organic layers were dried over anhydrous sodium sulfate and concentrated to dryness. The residue was purified by flash chromatography (40% EtOAc/Hexane) to give 0.61 g (73%) of the desired product. Analyzed by HPLC/MS.

25 Calculated weight 212.29. Found 213.15 (M+H).

Step 5 and 6:

A mixture of [S-(R*,R*)]- -butyl- -(propen-2-yl)- -hydroxymethyl- -butyrolactone (0.61 g, 2.88 mmol), NaOH (0.4 g, 10 mmol), THF (36 ml) , and water (9 ml) was stirred vigorously for 21 h at room temperature. Additional NaOH (0.12 g, 3 mmol) was added after 20 h. The mixture was concentrated and the residue was taken up into MeOH (40 ml). A solution of sodium periodate (2.4 g, 11.2 mmol) and H₂O (6 ml) was added and the mixture was stirred for 16 h at room temperature. After concentration, the aqueous residue was acidified by 1N HCl and extracted with DCM (2 x 30 ml). The combined organic layers was dried over anhydrous sodium sulfate, filtered, and concentrated to give 550 mg of the title compound. Analyzed by HPLC/MS. Calculated weight 198.26. Found 197.4

(M-H).

Step 7:

NaOMe (6.7 ml, 0.5 M in MeOH) was added to a solution of L-valine pyrrolidine amide HCl salt (0.69 g, 3.33 mmol) in MeOH (5 ml). The resulting solution was then added to [S-(R*,R*)]-butyl-(propen-2-yl)-hydroxy-butyrolactone (0.55 g, 2.78 mmol) in MeOH (10 ml). Sodium cyanoborohydride (0.24 g, 3.75 mmol) was added and the solution was stirred for 16 h at room temperature. After concentration, saturated aqueous ammonium chloride solution was added to the residue and extracted with DCM (2 x 30 ml). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated to provide 0.96 g of the uncyclized intermediate as oil. Analyzed by HPLC/MS. Calculated weight 352.52. Found 351.40 (M-H).

Step 8:

The uncyclized intermediate (0.96 g, 2.73 mmol) in THF (5 ml) was added to a solution of carbonyldiimidazole (0.53 g, 3.27 mmol) in THF (15 ml) and stirred for 16 h at room temperature. After concentration, the residue was taken up in EtOAc (30 ml) and washed with 1N HCl (20 ml) and brine (20 ml). The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated. The crude was purified by flash chromatography (75% EtOAc/Hexane) to provide 0.53 g (58 % yield) of the title compound. Analyzed by HPLC/MS. Calculated weight 334.51. Found 335.40 (M+H).


Step 9: Synthesis of compound 310

Ruthenium (IV) oxide hydrate (41 mg, 0.3 mmol) was added to a mixture of sodium periodate (2.55 g, 12 mmol), H₂O (33 ml), CH₃CN (33 ml) and CCl₄ (33 ml). The mixture was stirred for 20 min at room temperature and then NaHCO₃ (3.15 g, 37.5 mmol) and H₂O (17 ml) were added. After stirring for additional 5 min, a solution of pyrrolidineacetamide (0.5 g, 1.5 mmol) in CH₃CN (12 ml) was added. After stirring for 10 min, the residue was poured into a mixture of H₂O (100 ml) and EtOAc (200 ml). The aqueous layer was removed and the organic layer was extracted with 10% aqueous Na₂CO₃ solution (2 x 20 ml). The combined basic layers were acidified with 6N HCl and extracted with DCM (3 x 70 ml). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated. The crude product was purified by flash chromatography (EtOAc) to provide 108 mg (20%) of compound 310. Analyzed by HPLC/MS. Calculated weight 352.48. Found 353.40 (M+H).

Step 10: Synthesis of compound 311

To a stirring solution of acid (50 mg, 0.14 mmol) and triethylamine (23 ml, 0.17 mmol) in THF (2 ml) was added isobutylchloroformate (20 ml, 0.16 mmol) at 0°C. After

stirring for 15 min at 0°C, TMSONH₂ (36 ml, 0.21 mmol) was added and allowed the resulting mixture to stir for additional 2 h at room temperature. EtOAc (5 ml) was added to the solution and extracted with saturated aqueous ammonium chloride. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The crude material was



Scheme XXVIII

purified by flash chromatography (10% MeOH/DCM) to provide 24 mg (50%) of compound 311. Analyzed by HPLC/MS. Calculated weight 367.49. Found 366.40 (M-H).

5.9.1 Selectivity Profiles

See section 4.9 for description of selectivity assays.

The selectivity of the inhibitors was calculated as the ratio of the IC₅₀ for the compound versus thermolysin (THERM) divided by the IC₅₀ for the compound versus peptide deformylase (Fe-PDF). The metalloproteases carboxypeptidase (CPEP), collagenase (COLL) and angiotensin converting enzyme (ACE) were also used in these

studies but the IC₅₀ for thermolysin (THERM) was generally the smallest for these enzymes.

For Fe-PDF, inhibition is given as percent inhibition and for THERM, CPEP, COLL and ACE the inhibition values are the inhibition relative to controls with no inhibitor
5 (enzymatic reaction without test compound divided by reaction with test compound).

Table 17
Selectivity Profiles

	COMPOUND NUMBER	SELECTIVITY INDEX	Fe PDF	THER M	CPEP	Coll	ACE
10	10	30000	100	0.94	0.4	0.34	0.58
	20	15000	100	1.11	0.06	0.15	0.11
	28	30000	100	1.14	0.79	0.83	0.84
	35	15000	100	0.98	0.52	0.47	0.01
15	42	30000	100	0.9	0.5	0.62	0.2
	53	60000	100	1.04	0.15	0.28	0.19
	62	30000	100	1.09	0.34	0.28	-0.16
	64	30000	100	1.07	0.92	0.33	-0.45
20	65	10000	100	1.09	0.93	0.41	1.1
	66	7500	100	1.1	0.96	0.26	0.81
	86		100	0.35	0.28	0.32	0.12
	89		100	0.27	0.74	0.25	-0.26
25	90	15000	100	0.82	0.81	0.53	0.75
	97		100	0.39	0.75	0.47	0.53
	99	15000	100	0.51	0.93	0.38	0.97
	100	30000	100	0.42	0.79	0.67	0.79
30	109	10000	100	1.02	0.4	0.4	-0.05
	111	10000	100	0.97	0.45	0.51	0.16
	124	3750	100	0.91	0.8	0.77	0.78
	72	15000	100	0.16	0.73	0.11	0.4
35	74	3750	100	1.07	0.93	0.37	1

	COMPOUND NUMBER	SELECTIVITY INDEX	Fe PDF	THER M	CPEP	Coll	ACE
5	218	6000	100	0.88	0.39	0.51	0.6
	219	15000	100	0.95	0.15	0.6	-0.31
	221	10000	100	1.01	0.27	0.49	0.51
	201	10000	100	0.82	0.64	0.7	1.11
10	193	15000	100	1.01	0.97	0.93	0.89
	238	7500	100	0.83	0.58	0.91	0.99
	143	1500	100	0.06	0.96	0.65	1.1
	146	850	100	0.03	1.06	0.03	0.96
15	164	1667	100	0.88	0.62	0.97	1.53
	214	30000	100	0.54	0.93	1	0.74
	179	30000	100	0.59	0.96	0.82	0.28
	193	30000	100	0.57	0.04	1.2	0.84
20	315	1000	100	0	0.99	0.65	0.95
	316	167	102.5	0.01	0.95	0.8	1.02
	241	47	100	0.97	1	0.89	0.69
	274	15000	100	1.02	1.14	1.01	1
25	277	2727	100	0.84	0.94	1.07	0.72
	279	189	100	0.95	1.03	1.03	1.15
	234	166	104.1	0.99	0.96	1	0.97
	311	1667	100	0.91	1.1	1.02	1.94


30

35


THE CLAIMS

We Claim:


- 5 1. A compound represented by Formulas I- IV:




Formula I



Formula II



Formula III



Formula IV

including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein:

Z is NHOH or OR_a wherein R_a is H, alkyl or a biocleavable moiety;


X is C=O or O=S=O;

10 Y is heteroalkyl, heterocyclic, or substituted derivatives thereof;

R₁ is C₁-C₁₀ alkyl, cycloalkyl, aryl, heterocyclic, heteroalkyl or a substituted derivative thereof;


- 5 R_2 and R_3 together represent a 4, 5, 6, or 7 membered heterocyclic ring optionally substituted with one of the following substituents -OH, -CH₂OH, -O-C(=O)-heterocyclic, -O-C(=O)-NH-aryl, -NH-C(=O)-aryl, -C(=O)-N(C₂-C₆ alkyl)₂, -NH-(C₁-C₆ alkyl)-heterocyclic, -NH-C(=O)-NH-alkyl, -NH-C(=O)-NH-aryl, -NH-C(=O)-NH-heterocyclic; or
- R_2 and R_4 together form a ring through a -CH₂-CH₂- linkage wherein R_3 is H, C₁-C₁₀ alkyl, aryl, heterocyclic, heteroalkyl or a substituted derivative thereof; or R_2 is methyl;
- 10 R_3 , as defined above forms a ring with R_2 , or represents H, or is alkyl, heteroalkyl, aryl, heterocyclic, and substituted derivatives thereof;
- R_3 when not in a ring with R_2 is alkyl, heteroalkyl, aryl, heterocyclic, and substituted derivatives thereof;
- R_4 when not in a ring with R_2 is H, C₁-C₁₀ alkyl, aryl, heterocyclic, heteroalkyl or a substituted derivative thereof; or is together with R_2 as defined above to form
- 15 a ring through a -CH₂-CH₂- linkage;
- R_5 is -H, -NO₂, -NH₂, -NH-C(=O)-H, -NH-C(=O)-CH₃, -NH-S(O₂)-CH₃, -CH₂-NH-alkyl, -CH₂-NH-heteroalkyl, -CH₂-NH-heterocycl, or substituted derivatives thereof;
- 20 R_6 is -H, -NO₂, -NH₂, -NH-C(=O)-H, -NH-C(=O)-CH₃, -NH-S(O₂)-CH₃, -CH₂-NH-(alkyl), -CH₂-NH-heteroalkyl, -CH₂-NH-heterocycl, or substituted derivatives thereof;
- R_7 or R_8 is -CHR₁₀-C(=O)-NH-OH;
- R_7 or R_8 when not -CHR-C(=O)-NH-OH is alkyl, heteroalkyl, heterocycl, alkylaryl, alkylheterocyclic, or substituted derivatives thereof;
- 25 R_9 is H, alkyl, heteroalkyl, heterocycl, alkylaryl, alkylheterocyclic, or substituted derivatives thereof; and,
- R_{10} is H, alkyl, heteroalkyl, heterocycl, alkylaryl, alkylheterocyclic, or substituted derivatives thereof.

2. A compound of the formula:




including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R' is H, and R is selected from the from the group consisting of 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 2-fluoro-phenyl, 3-fluoro-phenyl, 4-fluoro-phenyl, 2-bromo-phenyl, 3-bromo-phenyl, phenyl, 4-methoxy-phenyl, 4-trifluoromethoxy-phenyl, 4-(N,N-Dimethyl-amino)-phenyl, 4-bromo-phenyl, 4-methyl-phenyl, 4-n-butyl-phenyl, 4-trifluoro-methyl-phenyl, 2-methoxy-phenyl, 2-ethyl-phenyl, 2-i-propyl-phenyl, 3-methyl-phenyl, 3-nitro-phenyl, 3-methyl-mercapto-phenyl, 4-chloro-3-nitro-phenyl, 4-chloro-3-trifluoromethyl-phenyl, 2,4-dichloro-phenyl, 2,4-dimethoxy-phenyl, 4-chloro-2-tri-fluoromethyl-phenyl, 2,5-difluoro-phenyl, 2-methoxy-5-chloro-phenyl, 2-methoxy-5-methyl-phenyl, 3,5-dimethyl-phenyl, 3,5-bis-(tri-fluoromethyl)-phenyl, 3,5-dichloro-phenyl, 4-(2,6-dichloro-pyridyl), n-propyl, benzyl, 2-(ethyl-2-thiophene), 4-phenoxy-phenyl, 4-trifluoro-methanemercapto-phenyl, 2-phenyl-phenyl, and 2-phenoxy-phenyl.

3. A compound of the formula:




including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R is selected from the from the group consisting of N-pyrrolidine, N-pyrrolidine-2-methanol, N-piperidine, N-homo-piperidine, N-morpholine, and N-3S-pyrrolidinol.

4. A compound of the formula:



- including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R is selected from the group consisting of phenyl, 3,5-di-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 4-methoxy-phenyl, 4-(N,N-dimethylamino)-phenyl, 4-trifluoro-methoxy-phenyl, and 4-fluoro-phenyl, and R' is H.


5. A compound of the formula:



- including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof wherein R is selected from the group consisting of N-isobutylamine, N-cyclopentylamine, N-cyclohexylamine, N-tetrahydro-furfurylamine, N-furfurylamine, N-thiophenmethylamine, N-benzylamine, 2-aminothiazole, N-dimethylamine, N-pyrrolidine, N-piperidine, N-homopiperidine, N-morpholine, 2R,6S-dimethyl-morpholine, and N-thiomorpholine.


15

6. A compound of the formula:




including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R is selected from the group consisting of N-isobutylamine, N-cyclopentylamine, N-cyclohexylamine, N-tetrahydro-furfurylamine, N-furfurylamine,
 5 N-2-thiophenemethylamine, N-methyl-3-pyridyl, N-benzylamine, 2-aminothiazole, N-dimethylamine, N-piperidine, N-homo-piperidine, N-morpholine, N-2,6-dimethyl-morpholine, N-thiomorpholine, and N-Methyl-N-piperazine.

7. A compound of the formula:




10 including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R is selected from the group consisting of N-isobutylamine, N-cyclopentylamine, N-cyclohexylamine, N-tetrahydro-furfurylamine, N-furfurylamine, N-thiophene-methylamine, N-2-methylpyridyl, N-benzylamine, N-2-aminothiazole, N-dimethylamine, N-pyrrolidine, N-piperidine, N-homopiperidine, N-morpholine, N-
 15 2R,6S-dimethylmorpholine, N-thiomorpholine, and N-methyl-N-piperazine.

8. A compound of the formula:




including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R represents is selected from the group consisting of N-isobutylamine, N-cyclopentylamine, N-furfurylamine, N-thiophene-methylamine,
 5 N-2-methyl-2-pyridyl, N-benzyl, N-2-aminothiazole, N,N-dimethylamine, N-pyrrolidine, N-piperidine, N-homopiperidine, N-morpholine, 2,6-dimethyl-morpholine, and N-thiomorpholine.

9. A compound of the formula:



10 including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R is selected from the group consisting of N-2-aminothiazole,
 N-dimethylamine, N-2-methylpyridyl, N-cyclopentylamine, and N-thiomorpholine.


10. A compound of the formula:



including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R_1 is selected from the group consisting of 3,4-difluorophenyl, phenyl and cyclohexyl; and R_2 is selected from the group consisting of N-2-aminothiazole and N-pyrrolidine.

5


11. A compound of the formula:




including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R is selected from the group consisting of valinyl-pyrrolidinamide, amino-valeryl-N,N-dimethylamide, phenylalanyl-N,N-dimethylamide, methionyl-N,N-dimethylamide, O-benzyl-serinyl-N,N-dimethylamide, O-benzyl-tyrosinyl-N,N-dimethylamide, amino-valeryl pyrrolidinamide, phenylalanyl pyrrolidinamide, methionyl pyrrolidinamide, O-benzyl-serinyl pyrrolidinamide, O-benzyl-tyrosinyl pyrrolidinamide, valinyl-piperidinamide, amino-valeryl-piperidinamide, phenylalanyl piperidinamide, methionyl-piperidinamide, O-benzyl-serinyl piperidinamide, O-benzyl-tyrosinyl-piperidinamide, valinyl-benzylamide, amino-valeryl-benzylamide, phenylalanyl-benzylamide, methionyl-benzylamide, O-benzyl-tyrosinyl-benzylamide, O-benzyl-tyrosinyl-benzylamide, phenylalanyl anilnamide, N-butylamine, N-pentylamine, N-pyrrolidine, N-piperidine, N-furfurylamine, N-adamantylamine, N-aniline, N-[4-(N-morpholine)]-aniline, N-[4-pentamethyloxy]-aniline, N-(4-fluoro)-aniline, N-(3-bromo)-aniline, N-(4-bromo)-aniline, N-(4-chloro)-aniline, N-(3,5-dimethoxy)-aniline, N-(3-phenoxy)-aniline, N-(4-phenoxy)-aniline, N-(4-N-acetamidyl) aniline, N-serinyldimethyl amide, lysinyl-pyrrolidinamide, glyciny-(2-acetohydroxamate)-pyrrolidinamide, glyciny-piperidinamide, lysinyl-dimethylamide, lysinyl-piperidinamide, glyciny-benzyl amide, lysinyl-benzyl amide, glyciny-4-methoxy-anilnamide, lysinyl-4-methoxy-anilnamide, glyciny-(2-acetohydroxamate)-4-methoxy anilnamide, glyciny-(2-propiohydroxamate)-4-methoxy anilnamide, glyciny-(4-fluoro)-anilnamide, N-propionate-(4-fluoro)-anilnamide, N-2-

- piperidinemethanol, N-4-piperidineethanol, (N-benzylethanolamine, N,N-diethanolamine, N-methyl-N-[(3,4-dimethoxy)-2-phenylethyl]-amine, N-2-(methylamino)ethanol, N-pyrrolidine-2-pyrrolidinamide, N-pyrrolidine-3-benzylamide, N-piperazine-N-pyrrolidinamide, N-piperazine-N-benzylamide, N-azetidine-3-benzylamide, 3-pyrrolidinol, 2-pyrrolidinol, 3-piperidinol, N-3-benzylamide-azetidine, N-piperidine-2-pyrrolidinamide and N-azetidine-(3-O-anilinamide).


12. A compound of the formula:




or



or




or



including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof.


13. A compound of the formula:



5 or a pharmaceutically acceptable salt thereof, wherein the R side chains are independently selected from the group consisting of H, NO₂, NHC(=O)CH₃, NHC(=O)H, and NHS(O₂)CH₃.


14. A compound of the formula:

10



15 including enantiomers and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R is selected from the group consisting of 1-naphthalene 1,2,3,4-tetrahydride, n-butyl, ethyl 2-phenyl, cyclopentyl, 2-methyltetrahydrofuran, and 4-chlorophenyl.


15. A compound of the formula:



including enantiomers, and mixtures thereof, or a pharmaceutically acceptable salt thereof, wherein R is selected from the group consisting of 1-naphthalene 1,2,3,4-tetrahydride, n-butyl, ethyl 2-phenyl, cyclopentyl, 2-methyltetrahydrofuran, and 4-chlorophenyl.

5


16. A compound of the formula:



including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof wherein z is selected from the group consisting of -OH and -NH-OH, M is 0 or 1; and R is selected from the group consisting of alkyl, aryl, alkylaryl, heterocycyl and substituted derivatives thereof.

10

17. A compound of the formula:




- including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof wherein B is selected from the group consisting of -NH-OH, N-isobutylamine, N-cyclopentylamine, N-furfurylamine, N-thiophene-methylamine, N-methyl-2-pyridine, N-benzyl, N-2-aminothiazole, N,N-dimethylamine, N-pyrrolidine, N-piperidine, N-homopiperidine, N-morpholine, N-2,6-dimethyl-morpholine, N-thiomorpholine, N-3-pyrrolidinol, N-2-pyrrolidinol, N-3-piperidinol, and N-2-

15


- pyrrolidine-methanol; R_8 is selected from the group consisting of H, $-CH_3$, $-CH_2C(=O)NH-OH$, $-CH_2CH(CH_3)_2$, $-(CH_2)_3CH_3$, $-CH(CH_3)_2$, $-CH_2(C_6H_5)$, $-CH_2CH_2SCH_3$, alkyl, aryl, alkylaryl, heteroalkyl, heterocycyl, and substituted derivatives thereof; R_9 is selected from the group consisting of H, $-CH_3$, $-CH_2C(=O)NH-OH$, $-CH_2CH(CH_3)_2$, $-(CH_2)_3CH_3$, $-CH(CH_3)_2$, $-CH_2(C_6H_5)$, $-CH_2CH_2SCH_3$, alkyl, aryl, alkylaryl, heteroalkyl, heterocycyl and substituted derivatives thereof; and R_{10} is selected from the group consisting of H, $-CH_3$, $-CH_2C(=O)NH-OH$, $-CH_2CH(CH_3)_2$, $-(CH_2)_3CH_3$, $-CH(CH_3)_2$, $-CH_2(C_6H_5)$, $-CH_2CH_2SCH_3$, alkyl, aryl, alkylaryl, heteroalkyl, heterocycyl and substituted derivatives thereof.

- 10 18. A compound of the formula:




- including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof wherein R_7 is selected from the group consisting of H, $-CH_3$, $-CH_2C(=O)NH-OH$, $-CH_2CH(CH_3)_2$, $-(CH_2)_3CH_3$, $-CH(CH_3)_2$, $-CH_2(C_6H_5)$, $-CH_2CH_2SCH_3$, alkyl, aryl, alkylaryl, heteroalkyl, heterocycyl, and substituted derivatives thereof; R_9 is selected from the group consisting of H, $-CH_3$, $-CH_2C(=O)NH-OH$, $-CH_2CH(CH_3)_2$, $-(CH_2)_3CH_3$, $-CH(CH_3)_2$, $-CH_2(C_6H_5)$, $-CH_2CH_2SCH_3$, alkyl, aryl, alkylaryl, heteroalkyl, heterocycyl and substituted derivatives thereof; and R_{10} is selected from the group consisting of H, $-CH_3$, $-CH_2C(=O)NH-OH$, $-CH_2CH(CH_3)_2$, $-(CH_2)_3CH_3$, $-CH(CH_3)_2$, $-CH_2(C_6H_5)$, $-CH_2CH_2SCH_3$, alkyl, aryl, alkylaryl, heteroalkyl, heterocycyl and substituted derivatives thereof.

19. A compound of the formula:



or




including all enantiomers, diastereomers and mixtures thereof, or a pharmaceutically acceptable salt thereof.

5

20. A compound represented by Formula XXIII:

including all enantiomers, diastereomers, and mixtures thereof, or a pharmaceutically acceptable salt thereof,



Formula XXIII

wherein

- 10 R_a is alkyl which may be substituted by aryl, halogenaryl, or cycloalkyl;
 R_b and R_c may be the same or different and each represents hydrogen, alkyl, heterocycl-alkyl, aryl, aryl-alkyl, cycloalkyl, or heterocyclic ring which may be substituted by alkyl, or R_b and R_c together with nitrogen from 5, 6, or 7 membered heterocyclic ring which may contain another heteroatom selected from nitrogen, sulfur, or oxygen; and
 15 W is selected from the group consisting of sulfur and ethylenyl.

21. The compound according to claim 20 having Formula XXIV:
wherein R_b and R_c are the same as defined in claim 20.

22. The compound according to claim 20 having Formula XXV:
5 wherein R_b and R_c are the same as defined in claim 20

23. The compound according to claim 20 having Formula XXVI:
wherein R_a is selected from the group consisting of aryl, halogenoaryl, and cycloalkyl, and
 R_b and R_c are the same as defined in claim 20.

10

24. The compound according to claim 21, wherein R_b and R_c may be the
same or different and each represents a hydrogen atom, isobutyl, methyl, cyclopentyl,
cyclohexyl, tetrahydrofurfuryl, furfuryl, 2-thienylmethyl, benzyl, 2-thiazolyl, or R_b and R_c
together with nitrogen form a heterocyclic ring which is selected from the group consisting
15 of 1-pyrrolidinyl, piperidino, homopiperidino, morpholino, 2,6-dimethylmorpholino, and
thiomorpholino.

25. The compound according to claim 22, wherein R_b and R_c may be the
same or different and each a hydrogen atom, isobutyl, methyl, cyclopentyl, cyclohexyl,
20 tetrahydrofurfuryl, furfuryl, 2-thienylmethyl, 3-pyridylmethyl, benzyl, 2-thiazolyl, or R_b and
 R_c together with nitrogen form a heterocyclic ring which is selected from the group
consisting of piperidino, morpholino, 2,6-dimethylmorpholino, thiomorpholino,
4-methyl-1-piperazinyl, and 1-pyrrolidinyl.

25 26. The compound according to claim 23, wherein R_a is selected from the
group consisting of 3,4-difluorophenyl, phenyl, and cyclohexyl, and R_b and R_c may be the
same or different and each represents hydrogen or 2-thiazolyl or R_b and R_c together with
nitrogen form 1-pyrrolidinyl.

30

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 April 2002 (11.04.2002)

PCT

(10) International Publication Number
WO 02/028829 A3

(51) International Patent Classification⁷: **C07D 233/54**,
403/12, 207/26, 207/12, 409/12, 211/42, 277/06, 417/12,
277/46, 217/26, 207/16, 307/52, 211/22, 207/14, 401/12,
405/12, 205/04, 307/14, 333/68, C07C 259/06, C07D
295/18, C07C 323/60, C07D 295/12, C07C 237/36, A61K
31/40, 31/44, 31/16, A61P 31/00, A01N 43/36, 43/40,
43/42, 47/18, C07K 5/06

2-11-6-706 Miyahara Yodogawa, Osaka 532-0003 (JP).
GLUCHOWSKI, Charles; 154 Coolspring Court,
Danville, CA 94506 (US).

(74) Agents: **POISSANT, Brian, M.** et al.; Pennie & Edmonds
LLP, 1155 Avenue of the Americas, New York, NY 10036
(US).

(21) International Application Number: PCT/US01/29926

(22) International Filing Date:
24 September 2001 (24.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/234,967 25 September 2000 (25.09.2000) US
09/761,850 18 January 2001 (18.01.2001) US

(71) Applicant: **QUESTCOR PHARMACEUTICALS, INC.**
[US/US]; 3260 Whipple Road, Union City, CA 94587-
1217 (US).

(72) Inventors: **CHONG, Lee**; 37469 Marsten Drive, Ne-
mark, CA 94560 (US). **FRECHETTE, Roger**; 40 Estate
Lane, Reading, MA 01867 (US). **SCOTT, Carole**; 6593
Flanders Drive, Newark, CA 94560 (US). **TESTER,**
Richard; 877 Heatherstone Way, Mountain View, CA
94040 (US). **SMITH, Whitney**; 1122 Richmond Street, El
Cerrito, CA 94530 (US). **CHIBA, Katsumi**; Enoki 33-94
Suita, Osaka 564-0053 (JP). **SAKAMOTO, Masatoshi**;

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

— with international search report

(88) Date of publication of the international search report:
24 December 2003

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/028829 A3

(54) Title: PEPTIDE DEFORMYLASE INHIBITORS

(57) Abstract: The invention is directed to a novel class of compounds, which inhibits peptide deformylase, pharmaceutical com-
positions containing compounds that inhibit peptide deformylase, and method of treating various infections.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/29926

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D233/54 C07D403/12 C07D207/26 C07D207/12 C07D409/12
C07D211/42 C07D277/06 C07D417/12 C07D277/46 C07D217/26
C07D207/16 C07D307/52 C07D211/22 C07D207/14 C07D401/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K A61P A01N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 59568 A (BRITISH BIOTECH PHARM ;HUNTER MICHAEL GEORGE (GB); SPAVOLD ZOE MAR) 25 November 1999 (1999-11-25) cited in the application the whole document claim 1; example 1 ---	1-12, 19-26
P,X	WO 01 38561 A (QUESTCOR PHARMACEUTICALS INC) 31 May 2001 (2001-05-31) claims 9-19 --- -/--	1-12, 19-26

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

14 August 2002

Date of mailing of the international search report

20. 11. 2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seitner, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/29926

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D405/12 C07D205/04 C07D307/14 C07D333/68 C07C259/06
C07D295/18 C07C323/60 C07D295/12 C07C237/36 A61K31/40
A61K31/44 A61K31/16 A61P31/00 A01N43/36 A01N43/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>"Antibiotic Actinonin. I to VIII"</p> <p>JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, CHEMICAL SOCIETY. LETCHWORTH, GB, vol. 9, no. 9, 1975, pages 819-860, XP002119637</p> <p>ISSN: 0300-922X</p> <p>cited in the application</p> <p>the whole document</p> <p>---</p> <p>-/--</p>	<p>1-12,</p> <p>19-26</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 August 2002

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seitner, I

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/29926

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A01N43/42 A01N47/18 C07K5/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TAMAKI KAZUHIKO ET AL: "Synthesis and structure-active relationships of gelatinase inhibitors derived from matlystatins" CHEMICAL AND PHARMACEUTICAL BULLETIN, PHARMACEUTICAL SOCIETY OF JAPAN. TOKYO, JP, vol. 43, no. 11, 1995, pages 1883-1893, XP002165817 ISSN: 0009-2363 the whole document	1-12, 19-26
X	US 5 712 300 A (JACOBSEN E JON) 27 January 1998 (1998-01-27) cited in the application column 18, line 14 - line 18 examples 2,23,30-34,39 --- -/--	1-12, 19-26

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the International filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the International filing date but later than the priority date claimed

"T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

14 August 2002

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seitner, I

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/29926

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 01 44179 A (JAIN RAKESH ;NI ZHI JIE (US); PATEL DINESH V (US); VERSICOR INC (U) 21 June 2001 (2001-06-21) examples 1-35 claims 47,48	1-12, 19-26
X	--- SATO, TSUTOMU ET AL: "YM-24074, a new peptide antibiotic. II. Structural elucidation." JOURNAL OF ANTIBIOTICS (1996), 49(8), 811-814, XP002209585 the whole document	1-12, 19-26
X	--- TAMAKI, KAZUHIKO ET AL: "Total synthesis and inhibitory activity against gelatinase B of YL-01869P" J. ANTIBIOT. (1995), 48(1), 87-8, XP002209586 the whole document	1-12, 19-26
X	--- PATENT ABSTRACTS OF JAPAN vol. 018, no. 236 (C-1196), 6 May 1994 (1994-05-06) & JP 06 025183 A (BANYU PHARMACEUT CO LTD), 1 February 1994 (1994-02-01) abstract	1-12, 19-26
X	--- PATENT ABSTRACTS OF JAPAN vol. 015, no. 388 (C-0872), 2 October 1991 (1991-10-02) & JP 03 157372 A (YAMANOUCHI PHARMACEUT CO LTD), 5 July 1991 (1991-07-05) abstract	1-12, 19-26
X	--- PATENT ABSTRACTS OF JAPAN vol. 015, no. 197 (C-0833), 21 May 1991 (1991-05-21) & JP 03 053891 A (MEIJI SEIKA KAISHA LTD), 7 March 1991 (1991-03-07) abstract	1-12, 19-26
X	--- US 5 840 939 A (SPAVOLD ZOE MARIE ET AL) 24 November 1998 (1998-11-24) examples 5-8 column 2, line 31 - line 51	1-12, 19-26
X	--- WO 95 06031 A (IMMUNEX CORP) 2 March 1995 (1995-03-02) page 2, line 28 - line 29 page 3 example 10	1-12, 19-26
	--- -/--	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/29926

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 39704 A (BRITISH BIOTECH PHARM ;DAVIES STEPHEN JOHN (GB); HUNTER MICHAEL GE) 12 August 1999 (1999-08-12) example 18 claim 1	1-12, 19-26
X	--- GHOSE A K ET AL: "DETERMINATION OF PHARMACOPHORIC GEOMETRY FOR COLLAGENASE INHIBITORS USING A NOVEL COMPUTATIONAL METHOD AND ITS VERIFICATION USING MOLECULAR DYNAMICS, NMR, AND X-RAY CRYSTALLOGRAPHY" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC, US, vol. 117, no. 16, 1995, pages 4671-4682, XP002051616 ISSN: 0002-7863 figure 10; example X	1-12, 19-26
X	--- PATENT ABSTRACTS OF JAPAN vol. 1997, no. 05, 30 May 1997 (1997-05-30) & JP 09 003094 A (MERCIAN CORP), 7 January 1997 (1997-01-07) abstract	1-12, 19-26
X	--- INAOKA Y ET AL: "PROPIOXATINS A AND B, NEW ENKEPHALINASE B INHIBITORS III. TOTAL SYNTHESIS OF PROPIOXATIN A" JOURNAL OF ANTIBIOTICS, JAPAN ANTIBIOTICS RESEARCH ASSOCIATION. TOKYO, JP, vol. 39, no. 10, October 1986 (1986-10), pages 1382-1385, XP000978947 ISSN: 0021-8820 the whole document	1-12, 19-26
X	--- INAOKA Y ET AL: "PROPIOXATINS A AND B, NEW ENKEPHALINASE B INHIBITORS. IV. CHARACTERIZATION OF THE ACTIVE SITE OF THE ENZYME USING SYNTHETIC PROPIOXATIN ANALOGUES" JOURNAL OF BIOCHEMISTRY, JAPANESE BIOCHEMICAL SOCIETY, TOKYO, JP, vol. 104, no. 5, November 1988 (1988-11), pages 706-711, XP000978993 ISSN: 0021-924X example 17 -----	1-12, 19-26

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/29926

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; It is covered by claims Nos.:
see further information sheet

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1 (partially); 10 (partially); 11 (partially); 12 (completely); 19 (completely); 20 (partially)

The initial phase of the search of invention 1 revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT).

For these reasons, it appears impossible to execute a meaningful search and/or to issue a complete search report over the whole breadth of the above mentioned claims.

The search and the report for those claims can only be considered as complete for compounds of the general formula
 $\text{HO-NH-C(=O)-CH}_2\text{-CR}_1\text{R}_4\text{-C(=O)-NR}_2\text{R}_3$ in which $\text{R}_1=\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_3$ and none of R_2 or R_3 represent hydrogen.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1 (partially); 2-12 (completely); 19 (partially); 20-26 (completely)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=NHOH$ and $R_1=alkyl$ or a substituted derivative thereof.

2. Claim : 1 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=NHOH$ and $R_1=cycloalkyl$ or a substituted derivative thereof.

3. Claim : 1 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=NHOH$ and $R_1=aryl$ or a substituted derivative thereof.

4. Claim : 1 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=NHOH$ and $R_1=heterocyclic$ or a substituted derivative thereof.

5. Claim : 1 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=NHOH$ and $R_1=heteroalkyl$ or a substituted derivative thereof.

6. Claim : 1 (partially); 19 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=ORa$ according to claim 1 and $R_1=C_1-C_{10}$ alkyl or a substituted derivative thereof.

7. Claim : 1 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=ORa$ according to claim 1 and $R_1=cycloalkyl$ or a substituted derivative thereof.

8. Claim : 1 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=ORa$ according to claim 1 and $R_1=aryl$ or a substituted

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

derivative thereof.

9. Claim : 1 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=ORa$ according to claim 1 and R_1 =heterocyclic or a substituted derivative thereof.

10. Claim : 1 (partially)

Compounds of the formula $Z-C(=O)-CH_2-C(R_1)-C(=O)-N$ in which $Z=ORa$ according to claim 1 and R_1 =heteroalkyl or a substituted derivative thereof.

11. Claims: 1 (partially); 13-16 (completely)

Compounds according to formula III of claim 1 as well as formulae of claim 13-16.

12. Claims: 1 (partially); 17, 18 (completely)

Compounds according to formula IV of claim 1 as well as formulae of claims 17 and 18.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/29926

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9959568 A	25-11-1999	AU 3942199 A	06-12-1999
		BR 9910488 A	09-01-2001
		CA 2332713 A	25-11-1999
		CN 1300210 T	20-06-2001
		EP 1079819 A	07-03-2001
		GB 2353708 A	07-03-2001
		HU 0102290 A	28-11-2001
		JP 2002515427 T	28-05-2002
		NO 20005782 A	15-01-2001
		PL 344438 A	05-11-2001
		TR 200003376 T	21-03-2001
		US 6441042 B	27-08-2002
WO 0138561 A	31-05-2001	AU 1929301 A	04-06-2001
US 5712300 A	27-01-1998	AU 707180 B	01-07-1999
		AU 2052597 A	22-09-1997
		BR 9707947 A	27-07-1999
		CN 1210517 A	10-03-1999
		EP 0898562 A	03-03-1999
		JP 2000506163 T	23-05-2000
		NO 984112 A	06-11-1998
		NZ 330922 A	28-01-2000
		PL 328693 A	15-02-1999
		RU 2168497 C	10-06-2001
		TW 448172 B	01-08-2001
		WO 9732846 A	12-09-1997
		ZA 9701902 A	07-09-1998
WO 0144179 A	21-06-2001	AU 2268301 A	25-06-2001
		AU 2268401 A	25-06-2001
		EP 1237862 A	11-09-2002
		WO 0144178 A	21-06-2001
JP 06025183 A	01-02-1994	NONE	
JP 03157372 A	05-07-1991	NONE	
JP 03053891 A	07-03-1991	JP 1938070 C	09-06-1995
		JP 6070019 B	07-09-1994
US 5840939 A	24-11-1998	AT 198319 T	15-01-2001
		AT 195723 T	15-09-2000
		AU 5343696 A	07-11-1996
		DE 69609957 D	28-09-2000
		DE 69609957 T	01-03-2001
		DE 69611360 D	01-02-2001
		DE 69611360 T	13-06-2001
		EP 0821666 A	04-02-1998
		EP 0821668 A	04-02-1998
		ES 2153959 T	16-03-2001
		ES 2150116 T	16-11-2000
		WO 9633161 A	24-10-1996
		WO 9633165 A	24-10-1996
		JP 11503746 T	30-03-1999
		JP 11503748 T	30-03-1999
		US 5908851 A	01-06-1999

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inte Application No
PCT/US 01/29926

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9506031 A	02-03-1995	AU 5030298 A AU 687436 B AU 7569494 A EP 0715619 A FI 960803 A JP 9503201 T NO 960723 A NZ 271893 A US 5594106 A US 5629285 A	05-03-1998 26-02-1998 21-03-1995 12-06-1996 22-04-1996 31-03-1997 23-02-1996 24-11-1997 14-01-1997 13-05-1997
WO 9939704 A	12-08-1999	AU 749699 B AU 2529299 A BR 9907689 A CA 2320476 A CN 1298299 T EP 1052984 A GB 2349884 A HU 0102901 A JP 2002502815 T NO 20003969 A PL 342296 A TR 200002311 T US 2002165167 A US 6423690 B	04-07-2002 23-08-1999 14-11-2000 12-08-1999 06-06-2001 22-11-2000 15-11-2000 28-12-2001 29-01-2002 28-09-2000 04-06-2001 21-11-2000 07-11-2002 23-07-2002
JP 09003094 A	07-01-1997	NONE	